arXiv:1111.1893v3 [math.NT] 2 Jul 2012

Stable models of Lubin-Tate curves with level three

Naoki Imai and Takahiro Tsushima

Abstract

We construct a stable formal model of a Lubin-Tate curve of level three, and study the action of a Weil group and a division algebra on its stable reduction. Further, we study a structure of cohomology of the Lubin-Tate curve. Our study is purely local and includes the case where the characteristic of the residue field of a local field is two.

Introduction

Let K be a non-archimedean local field with a finite residue field k of characteristic p. Let $\mathfrak{p}$ be the maximal ideal of $\mathcal{O}_K$. Let n be a natural number. We write $\mathrm{LT}(\mathfrak{p}^n)$ for the Lubin-Tate curve with full level n over $\widehat{K}^{\mathrm{ur}}$. Let D be the central division algebra over K of invariant $1/2$. Let ℓ be a prime number different from p. Then the groups W_K, $GL_2(K)$ and $D^\times$ act on $\varinjlim_m H^1_{\mathrm{c}}(\mathrm{LT}(\mathfrak{p}^m)_{\widehat{K}^{\mathrm{ac}}}, \overline{\mathbb{Q}}_\ell)$, and these actions partially realize the local Langlands correspondence and the local Jacquet-Langlands correspondence for GL_2. The realization of the local Langlands correspondence was proved by global automorphic methods. However there is no known proof using only a local geometric method.

We put

$$K_1(\mathfrak{p}^n) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathcal{O}_K) \;\middle|\; c \equiv 0,\ d \equiv 1 \bmod \mathfrak{p}^n \right\}.$$

Let $\mathrm{LT}_1(\mathfrak{p}^n)$ be the Lubin-Tate curve with level $K_1(\mathfrak{p}^n)$ over $\widehat{K}^{\mathrm{ur}}$. Then the cohomology group

$$H^1_{\mathrm{c}}(\mathrm{LT}_1(\mathfrak{p}^n)_{\widehat{K}^{\mathrm{ac}}}, \overline{\mathbb{Q}}_\ell) = \Big(\varinjlim_m H^1_{\mathrm{c}}(\mathrm{LT}(\mathfrak{p}^m)_{\widehat{K}^{\mathrm{ac}}}, \overline{\mathbb{Q}}_\ell) \Big)^{K_1(\mathfrak{p}^n)}$$

will give representations of W_K and $D^\times$ that correspond to smooth irreducible representations of $GL_2(K)$ with conductor less than or equal to n. The purpose of this paper is to study this cohomology in the case $n = 3$. We note that 3 is the smallest integer which is a conductor of a primitive two-dimensional Galois representation. Our method is purely local and geometric. In fact, we construct a stable model of the connected Lubin-Tate curve $\mathbf{X}_1(\mathfrak{p}^3)$ with level $K_1(\mathfrak{p}^3)$ by using the theory of semi-stable coverings (cf. [CM, Section 2.3]). Our study includes the case where $p = 2$, and in this case, primitive Galois representations of conductor 3 appear in the cohomology of $\mathbf{X}_1(\mathfrak{p}^3)$. It gives geometric understanding of a realization of the primitive Galois representations.

Our method of the calculation of the stable reduction is similar to that in [CM]. In [CM], Coleman-McMurdy calculate the stable reduction of the modular curve $X_0(p^3)$ under the assumption $p \geq 13$. The calculation of the stable reductions in the modular curve setting is equivalent to that in the Lubin-Tate setting where $K = \mathbb{Q}_p$. As for the calculation of the stable reduction of the modular curve $X_1(p^n)$, it is given in [DR] if $n = 1$.

We explain the contents of this paper. In Section 1, we recall a definition of the connected Lubin-Tate curve, and study the action of a division algebra in a general setting. In Section 2, we study cohomology of Lubin-Tate curves as representation of $GL_2(K)$. By this result, we can calculate the genus of some Lubin-Tate curve. In Section 3, we construct a stable covering of the connected Lubin-Tate curve of level $K_1(\mathfrak{p}^2)$, which is used to study a covering of $\mathbf{X}_1(\mathfrak{p}^3)$.

In Section 4, we define several affinoid subspaces $\mathbf{Y}_{1,2}$, $\mathbf{Y}_{2,1}$ and $\mathbf{Z}^0_{1,1}$ of $\mathbf{X}_1(\mathfrak{p}^3)$, and calculate their reductions. We put $q = |k|$ and

$$\mathcal{S}_1 = \begin{cases} \mu_{2(q^2-1)}(k^{\mathrm{ac}}) & \text{if } q \text{ is odd,} \\ \mu_{q^2-1}(k^{\mathrm{ac}}) & \text{if } q \text{ is even.} \end{cases}$$

2010 *Mathematics Subject Classification.* Primary: 11G20; Secondary: 11F80.

The reductions of $\mathbf{Y}_{1,2}$ and $\mathbf{Y}_{2,1}$ are defined by $x^q y - xy^q = 1$. This affine curve has genus $q(q-1)/2$, and is called the Deligne-Lusztig curve for $SL_2(\mathbb{F}_q)$ or the Drinfeld curve. The reduction $\overline{\mathbf{Z}}^0_{1,1}$ of $\mathbf{Z}^0_{1,1}$ is defined by $Z^q + X^{q^2-1} + X^{-(q^2-1)} = 0$. This affine curve has genus 0 and singularities at $X \in \mathcal{S}_1$.

Next, we analyze tubular neighbourhoods $\{\mathcal{D}_\zeta\}_{\zeta \in \mathcal{S}_1}$ of the singular points of $\overline{\mathbf{Z}}^0_{1,1}$. If q is odd, $\mathcal{D}_\zeta$ is a basic wide open space with the underlying affinoid $\mathbf{X}_\zeta$. The reduction of $\mathbf{X}_\zeta$ is the Artin-Schreier curve of degree 2 defined by $z^q - z = w^2$. This affine curve has genus $(q-1)/2$.

On the other hand, if q is even, it is harder to analyze $\mathcal{D}_\zeta$, because the space $\mathcal{D}_\zeta$ is not basic. First, we find an affinoid $\mathbf{P}^0_\zeta$. The reduction $\overline{\mathbf{P}}^0_\zeta$ of $\mathbf{P}^0_\zeta$ has genus 0 and singular points parametrized by $\zeta' \in k^\times$. Secondly, we analyze the tubular neighborhoods of singular points of $\overline{\mathbf{P}}^0_\zeta$. As a result, we find an affinoid $\mathbf{X}_{\zeta,\zeta'}$, whose reduction $\overline{\mathbf{X}}_{\zeta,\zeta'}$ is defined by $z^2 + z = w^3$. The smooth compactification of this curve is the unique supersingular elliptic curve over k^{ac}, whose j-invariant is 0, and its cohomology gives a primitive Galois representation. By using these affinoid spaces, we construct a covering $\mathcal{C}_1(\mathfrak{p}^3)$ of $\mathbf{X}_1(\mathfrak{p}^3)$.

In Section 5, we calculate the action of the division algebra $\mathcal{O}_D^\times$ on the reductions of the affinoid spaces in $\mathbf{X}_1(\mathfrak{p}^3)$. In Section 6, we calculate an action of a Weil group on the reductions. We construct an $SL_2(\mathbb{F}_3)$-Galois extension of K^{ur}, and show that the Weil action on $\overline{\mathbf{X}}_{\zeta,\zeta'}$ up to translations factors through the Weil group of the constructed extension. For such a Galois extension, see also [We, 31].

In Section 7, we show that the covering $\mathcal{C}_1(\mathfrak{p}^3)$ is semi-stable. To show this, we calculate the summation of the genera of the reductions of the affinoid spaces in $\mathbf{X}_1(\mathfrak{p}^3)$, and compare it with the genus of $\mathbf{X}_1(\mathfrak{p}^3)$. Using the the constructed semi-stable model, we study a structure of cohomology of $\mathbf{X}_1(\mathfrak{p}^3)$.

The dual graph of the semi-stable reduction of $\mathbf{X}_1(\mathfrak{p}^3)$ in the case where q is even is the following:


$\overline{\mathbf{Y}}^{\mathrm{c}}_{1,2}$
$\overline{\mathbf{Z}}^{0,\mathrm{c}}_{1,1}$
$\overline{\mathbf{Y}}^{\mathrm{c}}_{2,1}$
$\overline{\mathbf{P}}^{0,\mathrm{c}}_{\zeta_1}$
$\overline{\mathbf{P}}^{0,\mathrm{c}}_{\zeta_{q^2-1}}$
$\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta_1,\zeta'_1}$
$\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta_1,\zeta'_{q-1}}$
$\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta_{q^2-1},\zeta'_1}$
$\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta_{q^2-1},\zeta'_{q-1}}$

where $\mu_{q^2-1}(k^{\mathrm{ac}}) = \{\zeta_1, \ldots, \zeta_{q^2-1}\}$, $k^\times = \{\zeta'_1, \ldots, \zeta'_{q-1}\}$ and X^{c} denotes the smooth compactification of X for a curve X over k^{ac}. The constructed semi-stable model is in fact stable, except in the case where $q = 2$. If $q = 2$, we get the stable model by blowing down some $\mathbb{P}^1$-components.

The realization of the local Jacquet-Langlands correspondence in cohomology of Lubin-Tate curves was proved in [M2] by a purely local method. Therefore, the remaining essential part of the study of the realization of the local Langlands correspondence is to study actions of Weil groups and division algebras. In the paper [IT2], we give a purely local proof of the realization of the local Langlands correspondence for representations of conductor three using the result of this paper.

Acknowledgment

The authors thank Seidai Yasuda for helpful discussion on the subject in Paragraph 6.2.2.

Notation

In this paper, we use the following notation. Let K be a non-archimedean local field. Let $\mathcal{O}_K$ denote the ring of integers and k the residue field of characteristic $p > 0$. We fix a uniformizer ϖ of K. Let $q = |k|$. For any finite extension F of K, let G_F denote the absolute Galois group of F, W_F denote the Weil group of F and I_F denote the inertia subgroup of W_F. We fix an algebraic closure K^{ac}. The completion of K^{ac} is denoted by $\mathbf{C}$. Let $\mathcal{O}_{\mathbf{C}}$ be the ring of integers of $\mathbf{C}$ and k^{ac} the residue field of $\mathbf{C}$. For an element $a \in \mathcal{O}_{\mathbf{C}}$, we write $\bar{a}$ for the image of a by the reduction map $\mathcal{O}_{\mathbf{C}} \to k^{\mathrm{ac}}$. Let $v(\cdot)$ denote a valuation of $\mathbf{C}$ such that $v(\varpi) = 1$. Let K^{ur} denote the maximal unramified extension of K in K^{ac}. The completion of K^{ur} is denoted by $\widehat{K}^{\mathrm{ur}}$. For $a, b \in \mathbf{C}$ and a rational number $\alpha \in \mathbb{Q}_{\geq 0}$, we write $a \equiv b \pmod{\alpha}$ if we have $v(a-b) \geq \alpha$, and $a \equiv b \pmod{\alpha+}$ if we have $v(a-b) > \alpha$. For a curve X over k^{ac}, we denote by X^c the smooth compactification of X, and the genus of X means the genus of X^{c}. For an affinoid $\mathbf{X}$, we write $\overline{\mathbf{X}}$ for its reduction. The category of sets is denoted by **Set**. For a representation τ of a group, the dual representation of τ is denoted by τ^*. We fix $\varpi_0 \in K^{\mathrm{ac}}$ such that $\varpi_0^{24q^4(q^2-1)} = \varpi$, and $\varpi^{m/(24q^4(q^2-1))}$ denotes ϖ_0^m for any integer m.

1 Preliminaries

1.1 The universal deformation

Let Σ denote a formal $\mathcal{O}_K$-module of dimension 1 and height 2 over k^{ac}, which is unique up to isomorphism. Let n be a natural number. We define $K_1(\mathfrak{p}^n)$ as in the introduction. In the following, we define the connected Lubin-Tate curve $\mathbf{X}_1(\mathfrak{p}^n)$ with level $K_1(\mathfrak{p}^n)$.

Let $\mathcal{C}$ be the category of Noetherian complete local $\mathcal{O}_{\widehat{K}^{\mathrm{ur}}}$-algebras with residue field k^{ac}. For $A \in \mathcal{C}$, a formal $\mathcal{O}_K$-module $\mathcal{F} = \operatorname{Spf} A[[X]]$ over A and an A-valued point P of $\mathcal{F}$, the corresponding element of maximal ideal of A is denoted by $x(P)$. We consider the functor

$$\mathcal{A}_1(\mathfrak{p}^n) \colon \mathcal{C} \to \mathbf{Set}; A \mapsto [(\mathcal{F}, \iota, P)],$$

where $\mathcal{F}$ is a formal $\mathcal{O}_K$-module over A with an isomorphism $\iota \colon \Sigma \simeq \mathcal{F} \otimes_A k^{\mathrm{ac}}$ and P is a ϖ^n-torsion point of $\mathcal{F}$ such that

$$\prod_{a \in \mathcal{O}_K/\varpi^n \mathcal{O}_K} \bigl(X - x([a]_{\mathcal{F}}(P))\bigr) \;\Big|\; [\varpi^n]_{\mathcal{F}}(X)$$

in $A[[X]]$. This functor is represented by a regular local ring $\mathcal{R}_1(\mathfrak{p}^n)$. We write $\mathfrak{X}_1(\mathfrak{p}^n)$ for $\operatorname{Spf} \mathcal{R}_1(\mathfrak{p}^n)$. Its generic fiber is denoted by $\mathbf{X}_1(\mathfrak{p}^n)$, which we call the connected Lubin-Tate curve with level $K_1(\mathfrak{p}^n)$. The space $\mathbf{X}_1(\mathfrak{p}^n)$ is a rigid analytic curve over $\widehat{K}^{\mathrm{ur}}$. We can define the Lubin-Tate curve $\mathrm{LT}_1(\mathfrak{p}^n)$ with level n by changing $\mathcal{C}$ to be the category of $\mathcal{O}_{\widehat{K}^{\mathrm{ur}}}$-algebras where ϖ is nilpotent, and ι to be a quasi-isogeny $\Sigma \otimes_{k^{\mathrm{ac}}} A/\varpi A \to \mathcal{F} \otimes_A A/\varpi A$. We consider $\mathrm{LT}_1(\mathfrak{p}^n)$ as a rigid analytic curve over $\widehat{K}^{\mathrm{ur}}$.

The ring $\mathcal{R}_1(1)$ is isomorphic to the ring of formal power series $\mathcal{O}_{\widehat{K}^{\mathrm{ur}}}[[u]]$. We simply write $\mathcal{B}(1)$ for $\operatorname{Spf} \mathcal{O}_{\widehat{K}^{\mathrm{ur}}}[[u]]$. Let $B(1)$ denote an open unit ball such that $B(1)(\mathbf{C}) = \{u \in \mathbf{C} \mid v(u) > 0\}$. The generic fiber of $\mathcal{B}(1)$ is equal to $B(1)$. Then, the space $\mathbf{X}_1(1)$ is identified with $B(1)$. Let $\mathcal{F}^{\mathrm{univ}}$ denote the universal formal $\mathcal{O}_K$-module over $\mathfrak{X}_1(1)$.

In this subsection, we choose a parametrization of $\mathfrak{X}_1(1) \simeq \mathcal{B}(1)$ such that the universal formal $\mathcal{O}_K$-module has a simple form. Let $\mathcal{F}$ be a formal $\mathcal{O}_K$-module of dimension 1 over a flat $\mathcal{O}_K$-algebra R. For an invariant differential ω on $\mathcal{F}$, a logarithm of $\mathcal{F}$ means a unique isomorphism $F \colon \mathcal{F} \xrightarrow{\sim} \mathbb{G}_a$ over $R \otimes K$ with $dF = \omega$ (cf. [GH, 3]). In the sequel, we always take an invariant differential ω on $\mathcal{F}$ so that a logarithm F has the following form;

$$F(X) = X + \sum_{i \geq 1} f_i X^{q^i} \text{ with } f_i \in R \otimes K.$$

Let $F(X) = \sum_{i \geq 0} f_i X^{q^i} \in K[[u, X]]$ be the universal logarithm over $\mathcal{O}_K[[u]]$. By [GH, (5.5), (12.3), Proposition 12.10], the coefficients $\{f_i\}_{i \geq 0}$ satisfy $f_0 = 1$ and $\varpi f_i = \sum_{0 \leq j \leq i-1} f_j v_{i-j}^{q^j}$ for $i \geq 1$, where $v_1 = u$, $v_2 = 1$ and $v_i = 0$ for $i \geq 3$. Hence, we have the followings;

$$f_0 = 1,\ f_1 = \frac{u}{\varpi},\ f_2 = \frac{1}{\varpi}\left(1 + \frac{u^{q+1}}{\varpi}\right),\ f_3 = \frac{1}{\varpi^2}\left(u + u^{q^2} + \frac{u^{q^2+q+1}}{\varpi}\right), \cdots . \tag{1.1}$$

By [GH, Proposition 5.7] or [Ha, 21.5], if we set

$$\mathcal{F}^{\mathrm{univ}}(X, Y) = F^{-1}(F(X) + F(Y)),\ [a]_{\mathcal{F}^{\mathrm{univ}}}(X) = F^{-1}(aF(X)) \tag{1.2}$$

for $a \in \mathcal{O}_K$, it is known that these power series have coefficients in $\mathcal{O}_K[[u]]$ and define the universal formal $\mathcal{O}_K$-module $\mathcal{F}^{\mathrm{univ}}$ over $\mathcal{O}_{\widehat{K}^{\mathrm{ur}}}[[u]]$ of dimension 1 and height 2 with logarithm $F(X)$. We have the following approximation formula for $[\varpi]_{\mathrm{u}}(X)$.

Lemma 1.1. *We have the following congruence;*

$$[\varpi]_{\mathcal{F}^{\mathrm{univ}}}(X) \equiv \varpi X + uX^q + X^{q^2} - \frac{u}{\varpi}\{(uX^q + X^{q^2})^q - u^q X^{q^2} - X^{q^3}\}$$
$$\mod (\varpi^3, \varpi^2 X^q, u\varpi X^q, \varpi X^{q^2}, X^{q^3+1}).$$

Proof. This follows from a direct computation using the relation $F([\varpi]_{\mathcal{F}^{\mathrm{univ}}}(X)) = \varpi F(X)$ and (1.1). □

In the sequel, $\mathcal{F}^{\mathrm{univ}}$ means the universal formal $\mathcal{O}_K$-module with the identification $\mathfrak{X}_1(1) \simeq \mathcal{B}(1)$ given by (1.2), and we simply write $[a]_{\mathrm{u}}$ for $[a]_{\mathcal{F}^{\mathrm{univ}}}$. The reduction of (1.2) gives a simple model of Σ such that

$$X +_{\Sigma} Y = X + Y,\ [\zeta]_{\Sigma}(X) = \bar{\zeta} X \text{ for } \zeta \in \mu_{q-1}(\mathcal{O}_K),\ [\varpi]_{\Sigma}(X) = X^{q^2}. \tag{1.3}$$

We put

$$\mathfrak{A}_n = \mathcal{O}_{\widehat{K}^{\mathrm{ur}}}[[u, X_n]]/\big([\varpi^n]_{\mathrm{u}}(X_n)/[\varpi^{n-1}]_{\mathrm{u}}(X_n)\big).$$

Then there is a natural identification

$$\mathfrak{X}_1(\mathfrak{p}^n) \simeq \operatorname{Spf} \mathfrak{A}_n \tag{1.4}$$

that is compatible with the identification $\mathfrak{X}_1(1) \simeq \mathcal{B}(1)$. The Lubin-Tate curve $\mathbf{X}_1(\mathfrak{p}^n)$ is identified with the generic fiber of the right hand side of (1.4). We set $X_i = [\varpi^{n-i}]_{\mathrm{u}}(X_n)$ for $1 \leq i \leq n-1$. We write $\mathfrak{X}(1)$ for $\mathfrak{X}_1(1)$.

1.2 Action of a division algebra on $\mathfrak{X}_1(\mathfrak{p}^n)$

Let D be the central division algebra over K of invariant $1/2$. We write $\mathcal{O}_D$ for the ring of integers of D. In this subsection, we recall the left action of $\mathcal{O}_D^{\times}$ on the space $\mathfrak{X}_1(\mathfrak{p}^n)$.

Let K_2 be the unramified quadratic extension of K. Let k_2 be the residue field of K_2, and $\sigma \in \mathrm{Gal}(K_2/K)$ be the non-trivial element. The ring $\mathcal{O}_D$ has the following description; $\mathcal{O}_D = \mathcal{O}_{K_2} \oplus \varphi \mathcal{O}_{K_2}$ with $\varphi^2 = \varpi$ and $a\varphi = \varphi a^{\sigma}$ for $a \in \mathcal{O}_{K_2}$. We define an action of $\mathcal{O}_D$ on Σ by $\zeta(X) = \bar{\zeta} X$ for $\zeta \in \mu_{q^2-1}(\mathcal{O}_{K_2})$ and $\varphi(X) = X^q$. Then this give an isomorphism $\mathcal{O}_D \simeq \mathrm{End}(\Sigma)$ by [GH, Proposition 13.10].

Let $d = d_1 + \varphi d_2 \in \mathcal{O}_D^{\times}$, where $d_1 \in \mathcal{O}_{K_2}^{\times}$ and $d_2 \in \mathcal{O}_{K_2}$. By the definition of the action of $\mathcal{O}_D$ on Σ, we have

$$d(X) \equiv \bar{d}_1 X + (\bar{d}_2 X)^q \bmod (X^{q^2}). \tag{1.5}$$

We take a lifting $\tilde{d}(X) \in \mathcal{O}_{K_2}[[X]]$ of $d(X) \in k_2[[X]]$. Let $\mathcal{F}_{\tilde{d}}$ be the formal $\mathcal{O}_K$-module defined by

$$\mathcal{F}_{\tilde{d}}(X, Y) = \tilde{d}\big(\mathcal{F}^{\mathrm{univ}}(\tilde{d}^{-1}(X), \tilde{d}^{-1}(Y))\big),\ [a]_{\mathcal{F}_{\tilde{d}}}(X) = \tilde{d}\big([a]_{\mathrm{u}}(\tilde{d}^{-1}(X))\big)$$

for $a \in \mathcal{O}_K$. Then, we have an isomorphism

$$\tilde{d} \colon \mathcal{F}^{\mathrm{univ}} \xrightarrow{\sim} \mathcal{F}_{\tilde{d}};\ (u, X) \mapsto (u, \tilde{d}(X)).$$

By [GH, Proposition 14.7], the formal $\mathcal{O}_K$-module $\mathcal{F}_{\tilde{d}}$ with

$$\Sigma \xrightarrow{d^{-1}} \Sigma \xrightarrow{\iota} \mathcal{F}^{\mathrm{univ}} \otimes k^{\mathrm{ac}} \xrightarrow{\tilde{d} \otimes k^{\mathrm{ac}}} \mathcal{F}_{\tilde{d}} \otimes k^{\mathrm{ac}}$$

gives a isomorphism

$$d \colon \mathfrak{X}(1) \to \mathfrak{X}(1), \tag{1.6}$$

which is independent of a choice of a lifting $\tilde{d}$, such that there is the unique isomorphism

$$j \colon d^{*}\mathcal{F}^{\mathrm{univ}} \xrightarrow{\sim} \mathcal{F}_{\tilde{d}};\ (u, X) \mapsto (u, j(X))$$

satisfying $j(X) \equiv X \bmod (\varpi, u)$, where $d^{*}\mathcal{F}^{\mathrm{univ}}$ denotes the pull-back of $\mathcal{F}^{\mathrm{univ}}$ over $\mathfrak{X}(1)$ by the map (1.6). Hence, we have

$$[\varpi]_{d^{*}\mathcal{F}^{\mathrm{univ}}}(j^{-1}(X)) = j^{-1}([\varpi]_{\mathcal{F}_{\tilde{d}}}(X)). \tag{1.7}$$

On the other hand, we have the following isomorphism;

$$d^{*}\mathcal{F}^{\mathrm{univ}} \xrightarrow{\sim} \mathcal{F}^{\mathrm{univ}};\ (u, X') \mapsto (d(u), X').$$

Furthermore, we consider the following isomorphism under the identification (1.4);

$$\psi_d \colon \mathfrak{X}_1(\mathfrak{p}^n) \longrightarrow \mathfrak{X}_1(\mathfrak{p}^n);\ (u, X_n) \mapsto \big(d(u), j^{-1}(\tilde{d}(X_n))\big), \tag{1.8}$$

which depends only on d as in [GH, Proposition 14.7]. We put $d^*(X) = j^{-1}(\tilde{d}(X))$. We define a left action of d on $\mathfrak{X}_1(\mathfrak{p}^n)$ by $[(\mathcal{F}, \iota, P)] \mapsto [(\mathcal{F}, \iota \circ d^{-1}, P)]$. Then this action coincides with ψ_d by the definition.

By (1.5), we have

$$\tilde{d}^{-1}(X) = d_1^{-1}X - d_1^{-(q+1)} d_2^q X^q \bmod (\varpi, X^{q^2}) \tag{1.9}$$

in $\mathcal{O}_{K_2}[[X]]$. We use the following lemma later to compute the $\mathcal{O}_D^\times$-action on the stable reduction of $\mathbf{X}_1(\mathfrak{p}^3)$.

Lemma 1.2. *We assume $v(u) = 1/(2q)$. Let $d = d_1 + \varphi d_2 \in \mathcal{O}_D^\times$. We set $u' = d(u)$. We change variables as $u = \varpi^{1/(2q)}\tilde{u}$ and $u' = \varpi^{1/(2q)}\tilde{u}'$. Then, we have the followings:*

$$u' \equiv d_1^{-(q-1)} u (1 + d_1^{-q} d_2 u) \bmod (\varpi, u^3), \tag{1.10}$$

$$j^{-1}(X) \equiv X + d_1^{-q} d_2 u X \bmod (\varpi, u^2 X, u X^2). \tag{1.11}$$

Proof. We set $d^{-1} = d'_1 + \varphi d'_2$. Then $d'_1 \equiv d_1^{-1}$, $d'_2 \equiv -d_1^{-(q+1)} d_2 \pmod 1$. First, we prove (1.10). If $v(u) = 1/(2q)$, a function $w(u)$ in [GH, (25.11)] is well-approximated by a function $\varpi u (\varpi + u^{q+1})^{-1}$. By [GH, (25.13)], we have

$$\frac{\varpi u'}{\varpi + u'^{q+1}} \equiv \frac{d_1'^q \varpi u (\varpi + u^{q+1})^{-1} + \varpi d_2'^q}{d'_2 \varpi u (\varpi + u^{q+1})^{-1} + d'_1} \equiv \frac{\varpi u (d_1 - d_2^q u^q)}{d_1^q (\varpi + u^{q+1}) - d_2 \varpi u} \quad (\text{mod } 1+).$$

Hence, we acquire the following by $u = \varpi^{1/(2q)}\tilde{u}$ and $u' = \varpi^{1/(2q)}\tilde{u}'$;

$$\frac{\tilde{u}'}{\tilde{u}'^{q+1} + \varpi^{\frac{q-1}{2q}}} \equiv \frac{\tilde{u}(d_1 - \varpi^{\frac{1}{2}} d_2^q \tilde{u}^q)}{d_1^q \tilde{u}^{q+1} + \varpi^{\frac{q-1}{2q}} d_1^q - \varpi^{\frac{1}{2}} d_2 \tilde{u}} \quad (\text{mod } (1/2)+). \tag{1.12}$$

By taking an inverse of the congruence (1.12), we obtain

$$(\tilde{u}' - d_1^{-(q-1)}\tilde{u})^q \equiv \varpi^{\frac{q-1}{2q}} \left(\frac{\tilde{u}' - d_1^{-(q-1)}\tilde{u}}{d_1^{-(q-1)}\tilde{u}\tilde{u}'} \right) + \varpi^{\frac{1}{2}} (d_1^{q-2} d_2^q \tilde{u}^{2q} - d_1^{-1} d_2) \quad (\text{mod } (1/2)+). \tag{1.13}$$

Now, we set $\tilde{u}' - d_1^{-(q-1)}\tilde{u} = \varpi^{1/(2q)} x$. By substituting this to (1.13) and dividing it by $\varpi^{1/2}$, we obtain

$$(x - d_1^{1-2q} d_2 \tilde{u}^2)^q \equiv d_1^{2q-2} \tilde{u}^{-2} (x - d_1^{1-2q} d_2 \tilde{u}^2) \quad (\text{mod } 0+).$$

Since x is an analytic function of $\tilde{u}$, a congruence $x \equiv d_1^{1-2q} d_2 \tilde{u}^2 \pmod{0+}$ must hold. Hence we have $\tilde{u}' \equiv d_1^{-(q-1)} \tilde{u} (1 + \varpi^{1/2q} d_1^{-q} d_2 \tilde{u}) \pmod{(1/(2q))+}$ using $\tilde{u}' - d_1^{q-1}\tilde{u} = \varpi^{1/(2q)} x$. This implies (1.10), because u' is an analytic function of u.

By Lemma 1.1, (1.7) and (1.9), we have $u' j^{-1}(X)^q \equiv j^{-1}(u d_1^{-(q-1)} X^q) \bmod (\varpi, X^{q^2})$. Hence, the assertion (1.11) follows from (1.10) and $j^{-1}(X) \equiv X \bmod (\varpi, u)$. □

2 Cohomology of Lubin-Tate curve

Let ℓ be a prime number different from p. We take an algebraic closure $\overline{\mathbb{Q}}_\ell$ of $\mathbb{Q}_\ell$. Let $\mathrm{LT}(\mathfrak{p}^n)$ be the Lubin-Tate curve with full level n over $\widehat{K}^{\mathrm{ur}}$ (cf. [Da, 3.2]). We put $H^i_{\mathrm{LT},\varpi} = \varinjlim_n H^i_{\mathrm{c}}\big((\mathrm{LT}(\mathfrak{p}^n)/\varpi^{\mathbb{Z}})_{\mathbf{C}}, \overline{\mathbb{Q}}_\ell\big)$ for a non-negative integer i. Then we can define an action of $GL_2(K) \times D^\times \times W_K$ on $H^i_{\mathrm{LT},\varpi}$ for a non-negative integer i (cf. [Da, 3.2, 3.3]).

We write $\mathrm{Irr}(D^\times, \overline{\mathbb{Q}}_\ell)$ for the set of isomorphism classes of irreducible smooth representations of $D^\times$ over $\overline{\mathbb{Q}}_\ell$, and $\mathrm{Disc}(GL_2(K), \overline{\mathbb{Q}}_\ell)$ for the set of isomorphism classes of irreducible discrete series representations of $GL_2(K)$ over $\overline{\mathbb{Q}}_\ell$. Let $\mathrm{JL}\colon \mathrm{Irr}(D^\times, \overline{\mathbb{Q}}_\ell) \to \mathrm{Disc}(GL_2(K), \overline{\mathbb{Q}}_\ell)$ be the local Jacquet-Langlands correspondence. We denote by LJ the inverse of JL. We write St for the Steinberg representation of $GL_2(K)$.

Proposition 2.1. *We have isomorphisms*

$$H^1_{\mathrm{LT},\varpi} \simeq \bigoplus_{\pi} \pi^{\oplus 2 \dim \mathrm{LJ}(\pi)} \oplus \bigoplus_{\chi} (\mathrm{St} \otimes (\chi \circ \det)), \quad H^2_{\mathrm{LT},\varpi} \simeq \bigoplus_{\chi} (\chi \circ \det)$$

as representations of $GL_2(K)$, where π runs through irreducible cuspidal representations of $GL_2(K)$ whose central character is trivial on ϖ, and χ runs through characters of $K^\times$ satisfying $\chi(\varpi^2) = 1$.

Proof. First, we show the second isomorphism. Let $\mathbf{X}(\mathfrak{p}^n)$ be the connected Lubin-Tate curve with full level n over $\widehat{K}^{\mathrm{ur}}$ (cf. [St3, 2.1]). We put $H^2_{\mathbf{X}} = \varinjlim_n H^2_{\mathrm{c}}(\mathbf{X}(\mathfrak{p}^n)_{\mathbf{C}}, \overline{\mathbb{Q}}_\ell)$ and $GL_2(K)^0 = \{g \in GL_2(K) \mid \det g \in \mathcal{O}_K^\times\}$. Then $GL_2(K)^0$ acts on $H^2_{\mathbf{X}}$. By [St3, Theorem 4.4 (i)], we have $H^2_{\mathbf{X}} \simeq \bigoplus_\chi (\chi \circ \det)$ as representations of $GL_2(\mathcal{O}_K)$, where χ runs through characters of $\mathcal{O}_K^\times$. Let H be the kernel of $GL_2(K)^0 \to \mathrm{Aut}(H^2_{\mathbf{X}})$. Then $H = SL_2(K)$, because a normal subgroup of $GL_2(K)^0$ containing $SL_2(\mathcal{O}_K)$ is $SL_2(K)$ by [De, Lemme 2.2.5 (iii)]. Hence, we have $H^2_{\mathbf{X}} \simeq \bigoplus_\chi (\chi \circ \det)$ as representations of $GL_2(K)^0$, where χ runs through characters of $\mathcal{O}_K^\times$. The second isomorphism follows from this, because $H^2_{\mathrm{LT},\varpi} \simeq \text{c-Ind}_{GL_2(K)^0}^{GL_2(K)/\varpi^{\mathbb{Z}}} H^2_{\mathbf{X}}$.

Next, we show the first isomorphism. By [M2, Definition 6.2 and Theorem 6.6], the cuspidal part of $H^1_{\mathrm{LT},\varpi}$ is $\bigoplus_\pi \pi^{\oplus 2 \dim \mathrm{LJ}(\pi)}$. Here, we note that the characteristic of a local field is assumed to be zero in [M2], but the same proof works in the equal characteristic case. By [Far2, Théorème 4.3] and the Faltings-Fargues isomorphism (cf. [Fal] and [FGL]), we see that the non-cuspidal part of $H^1_{\mathrm{LT},\varpi}$ is the Zelvinsky dual of $H^2_{\mathrm{LT},\varpi}$. Therefore, we have the first isomorphism. □

3 Stable covering of Lubin-Tate curve with level two

In this section, we construct a stable covering of $\mathbf{X}_1(\mathfrak{p}^2)$. Let (u, X_2) be the parameter of $\mathbf{X}_1(\mathfrak{p}^2)$ given by the identification (1.4).

Let $\mathbf{Y}_{1,1}$, $\mathbf{W}_0$, $\mathbf{W}_{k^\times}$, $\mathbf{W}_{\infty,1}$, $\mathbf{W}_{\infty,2}$ and $\mathbf{W}_{\infty,3}$ be subspaces of $\mathbf{X}_1(\mathfrak{p}^2)$ defined by the following conditions respectively:

$$\begin{aligned}
\mathbf{Y}_{1,1}\colon & v(u) = \frac{1}{q+1},\ v(X_1) = \frac{q}{q^2-1},\ v(X_2) = \frac{1}{q(q^2-1)}.\\
\mathbf{W}_0\colon & 0 < v(u) < \frac{1}{q+1},\ v(X_1) = \frac{1-v(u)}{q-1},\ v(X_2) = \frac{1-qv(u)}{q(q-1)}.\\
\mathbf{W}_{k^\times}\colon & 0 < v(u) < \frac{1}{q+1},\ v(X_1) = \frac{1-v(u)}{q-1},\ v(X_2) = \frac{v(u)}{q(q-1)}.\\
\mathbf{W}_{\infty,1}\colon & 0 < v(u) < \frac{q}{q+1},\ v(X_1) = \frac{v(u)}{q(q-1)},\ v(X_2) = \frac{v(u)}{q^3(q-1)}.\\
\mathbf{W}_{\infty,2}\colon & v(u) \geq \frac{q}{q+1},\ v(X_1) = \frac{1}{q^2-1},\ v(X_2) = \frac{1}{q^2(q^2-1)}.\\
\mathbf{W}_{\infty,3}\colon & \frac{1}{q+1} < v(u) < \frac{q}{q+1},\ v(X_1) = \frac{1-v(u)}{q-1},\ v(X_2) = \frac{1-v(u)}{q^2(q-1)}.
\end{aligned}$$

We put $\mathbf{W}_\infty = \mathbf{W}_{\infty,1} \cup \mathbf{W}_{\infty,2} \cup \mathbf{W}_{\infty,3}$. Note that we have $\mathbf{X}_1(\mathfrak{p}^2) = \mathbf{Y}_{1,1} \cup \mathbf{W}_0 \cup \mathbf{W}^1_{k^\times} \cup \mathbf{W}_\infty$.

Proposition 3.1. *The Lubin-Tate curve $\mathbf{X}_1(\mathfrak{p}^2)$ is a basic wide open space with underlying affinoid $\mathbf{Y}_{1,1}$. Further, $\mathbf{W}_0$ and $\mathbf{W}_\infty$ are open annuli, and $\mathbf{W}_{k^\times}$ is a disjoint union of $q-1$ open annuli.*

Proof. We can calculate a semi-stable model of $\mathbf{X}_1(\mathfrak{p})$ from a good model $\mathcal{X}_1(1)$ of $\mathbf{X}_1(\mathfrak{p})$ by the similar argument as in this paper. In fact, we can show that $\mathbf{X}_1(\mathfrak{p})$ is isomorphic to an open annulus. Then we can prove the claim from the semi-stable model of $\mathbf{X}_1(\mathfrak{p})$ again by the similar argument. We omit the detail (cf. a proof of Theorem 7.14).

There is also a proof by direct calculations, which does not need cohomological arguments (cf. [IT3]). □

4 Reductions of affinoid spaces in $\mathbf{X}_1(\mathfrak{p}^3)$

4.1 Definitions of several subspaces in $\mathbf{X}_1(\mathfrak{p}^3)$

In this subsection, we define several subspaces of $\mathbf{X}_1(\mathfrak{p}^3)$. Let (u, X_3) be the parameter of $\mathbf{X}_1(\mathfrak{p}^3)$ given by the identification (1.4).

Let $\mathbf{Y}_{1,2}$, $\mathbf{Y}_{2,1}$ and $\mathbf{Z}^0_{1,1}$ be subspaces of $\mathbf{X}_1(\mathfrak{p}^3)$ defined by the following conditions respectively:

$$\begin{aligned}
&\mathbf{Y}_{1,2}\colon v(u)=\frac{1}{q+1},\ v(X_1)=\frac{q}{q^2-1},\ v(X_2)=\frac{1}{q(q^2-1)},\ v(X_3)=\frac{1}{q^3(q^2-1)}.\\
&\mathbf{Y}_{2,1}\colon v(u)=\frac{1}{q(q+1)},\ v(X_1)=\frac{q^2+q-1}{q(q^2-1)},\ v(X_2)=\frac{1}{q^2-1},\ v(X_3)=\frac{1}{q^2(q^2-1)}.\\
&\mathbf{Z}^0_{1,1}\colon v(u)=\frac{1}{2q},\ v(X_1)=\frac{2q-1}{2q(q-1)},\ v(X_2)=\frac{1}{2q(q-1)},\ v(X_3)=\frac{1}{2q^3(q-1)}.
\end{aligned}$$

We write down the following possible cases for (u,X_1,X_2);

$$\begin{aligned}
&1.\ 0<v(u)<\frac{1}{q+1},\ v(X_1)=\frac{1-v(u)}{q-1},\ v(X_2)=\frac{1-qv(u)}{q(q-1)},\\
&2.\ 0<v(u)<\frac{1}{q+1},\ v(X_1)=\frac{1-v(u)}{q-1},\ v(X_2)=\frac{v(u)}{q(q-1)},\\
&3.\ v(u)=\frac{1}{q+1},\ v(X_1)=\frac{q}{q^2-1},\ v(X_2)=\frac{1}{q(q^2-1)},\\
&4.\ 0<v(u)<\frac{q}{q+1},\ v(X_1)=\frac{v(u)}{q(q-1)},\ v(X_2)=\frac{v(u)}{q^3(q-1)},\\
&5.\ v(u)\geq\frac{q}{q+1},\ v(X_1)=\frac{1}{q^2-1},\ v(X_2)=\frac{1}{q^2(q^2-1)},\\
&6.\ \frac{1}{q+1}<v(u)<\frac{q}{q+1},\ v(X_1)=\frac{1-v(u)}{q-1},\ v(X_2)=\frac{1-v(u)}{q^2(q-1)}.
\end{aligned}\tag{4.1}$$

Next, we consider the following possible cases for (X_2,X_3);

$$\begin{aligned}
&1'.\ v(X_3^{q^2})=v(X_2)<v(uX_3^q),\ 2'.\ v(uX_3^q)=v(X_2)<v(X_3^{q^2}),\\
&3'.\ v(X_2)>v(X_3^{q^2})=v(uX_3^q),\ 4'.\ v(X_2)=v(X_3^{q^2})=v(uX_3^q).
\end{aligned}\tag{4.2}$$

Lemma 4.1. *For $2\leq i\leq 6$ in* (4.1) *and $2'\leq j'\leq 4'$ in* (4.2)*, the case i and j' does not happen.*

Proof. This is an easy exercise. □

Let $\mathbf{W}_{i,j'}$ be the subspace of $\mathbf{X}_1(\mathfrak{p}^3)$ defined by the conditions $1\leq i\leq 6$ in (4.1) and $1'\leq j'\leq 4'$ in (4.2). We note that $\mathbf{W}_{3,1'}=\mathbf{Y}_{1,2}$ and $\mathbf{W}_{1,4'}=\mathbf{Y}_{2,1}$. Let $\mathbf{W}^+_{1,1'}$, $\mathbf{W}^-_{1,1'}$ be the subspaces of $\mathbf{W}_{1,1'}$ defined by $1/(2q)<v(u)<1/(q+1)$, $1/(q(q+1))<v(u)<1/(2q)$ respectively.

4.2 Reductions of the affinoid spaces $\mathbf{Y}_{1,2}$ and $\mathbf{Y}_{2,1}$

In this subsection, we compute the reduction of the affinoid spaces $\mathbf{Y}_{1,2}$ and $\mathbf{Y}_{2,1}$. The reduction of $\mathbf{Y}_{2,1}$ and $\mathbf{Y}_{1,2}$ are defined by $x^qy-xy^q=1$. These curves have genus $q(q-1)/2$.

Proposition 4.2. *The reduction of* $\mathbf{Y}_{1,2}$ *is defined by* $x^qy-xy^q=1$.

Proof. We change variables as $u=\varpi^{1/(q+1)}\tilde{u}$, $X_1=\varpi^{q/(q^2-1)}x_1$, $X_2=\varpi^{1/(q(q^2-1))}x_2$ and $X_3=\varpi^{1/(q^3(q^2-1))}x_3$. By Lemma 1.1, we have

$$\tilde{u}\equiv -x_1^{-(q-1)},\ x_1\equiv\tilde{u}x_2^q+x_2^{q^2},\ x_2\equiv x_3^{q^2}\pmod{0+}.\tag{4.3}$$

Then we have $\tilde{u}=-x_1^{-(q-1)}+F_0(\tilde{u},x_1)$ for some function $F_0(\tilde{u},x_1)$ satisfying $v(F_0(\tilde{u},x_1))>v(\tilde{u})$. Substituting $\tilde{u}=-x_1^{-(q-1)}+F_0(\tilde{u},x_1)$ to $F_0(\tilde{u},x_1)$ and repeating it, we see that $\tilde{u}$ is written as a function of x_1. Similarly, by $x_2\equiv x_3^{q^2}$ (mod 0+), we can see that x_2 is written as a function of x_1 and x_3. By (4.3), we acquire

$$1\equiv\frac{x_3^{q^4}}{x_1}-\frac{x_3^{q^3}}{x_1^q}\pmod{0+}.\tag{4.4}$$

By setting $1+x_1^{-1}x_3^{q^2}=x_3^{q^3}t_1^{-1}$ and substituting this to (4.4), we obtain $t_1^q\equiv x_1$ (mod 0+) and hence $(1+x_3^qt_1^{-1})^q\equiv x_3^{q^3}t_1^{-1}$ (mod 0+). By setting $1+x_3^qt_1^{-1}=x_3^{q^2}t_2^{-1}$, we obtain $t_2^q\equiv t_1$ (mod 0+). Hence

$(1+x_3t_2^{-1})^q \equiv x_3^{q^2}t_2^{-1}$ (mod 0+). Finally, by setting $x=x_3$ and $1+x_3t_2^{-1}=x_3^q y$, we acquire $y^q \equiv t_2^{-1}$ (mod 0+). Hence we have $x^q y - xy^q \equiv 1$ (mod 0+). Note that

$$x=x_3,\ y=\frac{x_1(1+x_3^{q(q^2-1)}+x_3^{(q+1)(q^2-1)})+x_3^{q^2}}{x_1x_3^{q^3+q^2-1}}. \tag{4.5}$$

□

We put $\gamma_i=\varpi^{(q-1)/(2q^i)}$ for $1\le i\le 4$. We choose an element c_0 such that $c_0^q-\gamma_1^2c_0+1=0$. Note that we have $c_0\equiv -1$ (mod 0+). Further, we choose a q-th root $c_0^{1/q}$ of c_0.

Proposition 4.3. *The reduction of the space* $\mathbf{Y}_{2,1}$ *is defined by* $x^qy-xy^q=1$.

Proof. We change variables as $u=\varpi^{1/(q(q+1))}\tilde{u}$, $X_1=\varpi^{(q^2+q-1)/(q(q^2-1))}x_1$, $X_2=\varpi^{1/(q^2-1)}x_2$, and $X_3=\varpi^{1/(q^2(q^2-1))}x_3$. By Lemma 1.1, we have

$$\tilde{u}\equiv -x_1^{-(q-1)} \quad (\mathrm{mod}\ \frac{q^2-1}{q^2}+), \tag{4.6}$$

$$x_1\equiv \tilde{u}x_2^q+\gamma_1^2(x_2^{q^2}+x_2) \quad (\mathrm{mod}\ \frac{q^2-1}{q^2}+), \tag{4.7}$$

$$x_2\equiv x_3^{q^2}+\tilde{u}x_3^q \quad (\mathrm{mod}\ \frac{q-1}{q^2}+). \tag{4.8}$$

By (4.6) and (4.8), we can see that $\tilde{u}$ is written as a function of x_1, and that x_2 is written as a function of x_1 and x_3. We define a parameter t by

$$\frac{x_2}{x_1}=c_0+\gamma_2^2\frac{x_2^q}{t}. \tag{4.9}$$

We note that $v(t)=0$. By considering $x_1^{-1}\times$ (4.7), we have

$$\left(\frac{x_2}{x_1}\right)^q+1-\gamma_1^2\frac{x_2}{x_1}\equiv\gamma_1^2\frac{x_2^{q^2}}{x_1} \quad (\mathrm{mod}\ \frac{q^2-1}{q^2}+). \tag{4.10}$$

By substituting (4.9) to the left hand side of the congruence (4.10), and dividing it by $\gamma_1^2x_2^{q^2}$, we acquire

$$x_1\equiv t^q\left(1-\gamma_2^2\frac{t^{q-1}}{x_2^{q(q-1)}}\right)^{-1} \quad (\mathrm{mod}\ \frac{q-1}{q^2}+). \tag{4.11}$$

By this congruence, we can see that x_1 is written as a function of t and x_3. By considering $x_1^{-1}\times$ (4.8), we acquire

$$c_0+\gamma_2^2\frac{x_2^q}{t}\equiv\frac{x_3^{q^2}}{x_1}-\left(\frac{x_3}{x_1}\right)^q \quad (\mathrm{mod}\ \frac{q-1}{q^2}+) \tag{4.12}$$

by (4.9). Substituting (4.11) to (4.12), we have

$$\left(c_0^{1/q}-\frac{x_3^q}{t}+\frac{x_3}{x_1}\right)^q\equiv-\gamma_2^2\frac{(x_2+x_3)^{q^2}}{tx_2^{q(q-1)}} \quad (\mathrm{mod}\ \frac{q-1}{q^2}+). \tag{4.13}$$

By (4.9) and $c_0\equiv -1$ (mod 0+), we have $x_2\equiv -x_1$ (mod 0+). Therefore, we acquire $(x_2+x_3)^q\equiv x_2^{q-1}x_3^{q^2}$ (mod 0+) by (4.6) and (4.8). In particular, we obtain $v(x_2+x_3)=0$. We introducing a new parameter t_1 as

$$c_0^{1/q}-\frac{x_3^q}{t}+\frac{x_3}{x_1}=-\gamma_3^q\frac{(x_2+x_3)^q}{t_1x_2^{q-1}}. \tag{4.14}$$

Substituting this to the left hand side of the congruence (4.13), and dividing it by $-\gamma_2^2x_2^{-q(q-1)}(x_2+x_3)^{q^2}$, we acquire $t\equiv t_1^q$ (mod 0+). By this congruence, we can see that t is written as a function of t_1 and x_3. By (4.14), we obtain $x_3\equiv t_1^{q^2}(1+x_3t_1^{-1})^q$ (mod 0+) using $t\equiv t_1^q$ (mod 0+) and $x_1\equiv t^q$ (mod 0+). Hence, by setting $x=t_1^{-1}$ and $y=t_1^q(1+x_3t_1^{-1})$, we acquire $x^qy-yx^q\equiv 1$ (mod 0+). □

4.3 Reduction of the affinoid space $\mathbf{Z}_{1,1}^0$

In this subsection, we calculate the reduction of the affinoid space $\mathbf{Z}_{1,1}^0$. We define $\mathcal{S}_1$ as in the introduction. The reduction $\overline{\mathbf{Z}}_{1,1}^0$ is defined by $Z^q + x_3^{q^2-1} + x_3^{-(q^2-1)} = 0$. This affine curve has genus 0 and singularities at $x_3 \in \mathcal{S}_1$.

We put $\omega_i = \varpi^{1/(2q^i(q-1))}$ and $\epsilon_i = 1/(2q^i)$ for $1 \leq i \leq 4$. We change variables as $x = \omega_1^{q-1}\tilde{u}$, $X_1 = \omega_1^{2q-1}x_1$, $X_2 = \omega_1 x_2$ and $X_3 = \omega_3 x_3$. By Lemma 1.1, we have

$$\tilde{u} \equiv -x_1^{-(q-1)} \pmod{\frac{1}{2}+}, \tag{4.15}$$

$$x_1 \equiv \tilde{u}x_2^q + \gamma_1 x_2^{q^2} + \gamma_1^2 x_2 \pmod{\frac{1}{2}+}, \tag{4.16}$$

$$x_2 \equiv x_3^{q^2} + \gamma_2 \tilde{u} x_3^q \pmod{\epsilon_1+}. \tag{4.17}$$

Note that we have $v(\gamma_1^2) > 1/2$ if $q \neq 2$. By (4.15) and (4.17), we can see that $\tilde{u}$ is written as a function of x_1, and that x_2 is written as a function of x_1 and x_3. We define a parameter t by

$$\frac{x_2}{x_1} = -1 + \gamma_2 \frac{x_2^q}{t}. \tag{4.18}$$

By considering $x_1^{-1} \times$ (4.16), we acquire

$$\left(\frac{x_2}{x_1} + 1\right)^q \equiv \gamma_1 \frac{x_2^{q^2}}{x_1}\left(1 + \frac{\gamma_1}{x_2^{q^2-1}}\right) \pmod{\frac{1}{2}+} \tag{4.19}$$

by (4.15). Substituting (4.18) to (4.19), and dividing it by $\gamma_1 x_2^{q^2}$, we obtain

$$x_1 \equiv t^q\left(1 + \frac{\gamma_1}{x_2^{q^2-1}}\right) \pmod{\epsilon_1+}. \tag{4.20}$$

Therefore we have $v(t) = 0$. By considering $x_1^{-1} \times$ (4.17), we acquire

$$\left(1 + \frac{x_3^q}{t}\right)^q - \gamma_1 \frac{x_3^{q^2}}{t^q x_2^{q^2-1}} \equiv \gamma_2\left(\frac{x_2^q}{t} + \left(\frac{x_3}{x_1}\right)^q\right) \pmod{\epsilon_1+} \tag{4.21}$$

by (4.15), (4.18) and (4.20). We define a parameter Z_0 by

$$1 + \frac{x_3^q}{t} = \gamma_3 Z_0. \tag{4.22}$$

We note that $v(Z_0) \geq 0$. Substituting this to (4.21), and dividing it by γ_2, we obtain

$$Z_0^q \equiv \frac{x_2^q}{t} + \left(\frac{x_3}{x_1}\right)^q + \gamma_2^{q-1}\frac{x_3^{q^2}}{t^q x_2^{q^2-1}} \pmod{\epsilon_2+}. \tag{4.23}$$

By (4.22) and (4.23), we acquire

$$\left(Z_0 + \frac{x_2}{x_3} - \frac{x_3}{x_1}\right)^q \equiv \gamma_3\left(\frac{x_2}{x_3}\right)^q Z_0 + \gamma_2^{q-1}\frac{x_3^{q^2}}{t^q x_2^{q^2-1}} \pmod{\epsilon_2+}. \tag{4.24}$$

We introduce a new parameter Z as

$$Z_0 + \frac{x_2}{x_3} - \frac{x_3}{x_1} = \gamma_4 \frac{x_2}{x_3} Z. \tag{4.25}$$

We note that $v(Z) \geq 0$. Substituting this to the left hand side of the congruence (4.24), and dividing it by $\gamma_3(x_2/x_3)^q$, we acquire

$$Z^q \equiv Z_0 + \gamma_3^{q^2-q-1}\frac{x_3^{q(q+1)}}{t^q x_2^{q^2+q-1}} \pmod{\epsilon_3+}. \tag{4.26}$$

By substituting (4.25) to (4.26), we obtain

$$Z^q + x_3^{q^2-1}(1 - \gamma_4 Z) + x_3^{-(q^2-1)} \equiv -\gamma_3^{q^2-q-1} x_3^{-q(q^2-1)(q+1)} \pmod{\epsilon_3+} \tag{4.27}$$

by (4.17), (4.20) and (4.22). Note that we have $v(\gamma_3^{q^2-q-1}) > \epsilon_3$, if $q \neq 2$.

Proposition 4.4. *The reduction of the space* $\mathbf{Z}^0_{1,1}$ *is defined by* $Z^q + x_3^{q^2-1} + x_3^{-(q^2-1)} = 0$. *This affine curve has genus* 0 *and singularities at* $x_3 \in \mathcal{S}_1$.

Proof. The required assertion follows from the congruence (4.27) modulo 0+. ☐

Definition 4.5. *1. For any* $\zeta \in \mathcal{S}_1$, *we define a subspace* $\mathcal{D}_\zeta \subset \mathbf{Z}^0_{1,1} \times_{\widehat{K}^{\mathrm{ur}}} \widehat{K}^{\mathrm{ur}}(\omega_3)$ *by* $\bar{x}_3 = \zeta$. *We call the space* $\mathcal{D}_\zeta$ *a singular residue class of* $\mathbf{Z}^0_{1,1}$.
2. We define a subspace $\mathbf{Z}_{1,1} \subset \mathbf{Z}^0_{1,1} \times_{\widehat{K}^{\mathrm{ur}}} \widehat{K}^{\mathrm{ur}}(\omega_3)$ *by the complement* $\mathbf{Z}^0_{1,1} \times_{\widehat{K}^{\mathrm{ur}}} \widehat{K}^{\mathrm{ur}}(\omega_3) \setminus \bigcup_{\zeta \in \mathcal{S}_1} \mathcal{D}_\zeta$.

Proposition 4.6. *The reduction of the space* $\mathbf{Z}_{1,1}$ *is defined by* $Z^q + x_3^{q^2-1} + x_3^{-(q^2-1)} = 0$ *with* $x_3 \notin \mathcal{S}_1$.

Proof. This follows from Proposition 4.4. ☐

4.4 Analysis of the singular residue classes of $\mathbf{Z}^0_{1,1}$

In this subsection, we analyze the singular residue classes $\{\mathcal{D}_\zeta\}_{\zeta \in \mathcal{S}_1}$ of $\mathbf{Z}^0_{1,1}$. If q is odd, the space $\mathcal{D}_\zeta$ is a basic wide open space with an underlying affinoid $\mathbf{X}_\zeta$, whose reduction $\overline{\mathbf{X}}_\zeta$ is defined by $z^q - z = w^2$. On the other hand, if q is even, the situation is slightly complicated, because the space $\mathcal{D}_\zeta$ is not basic wide open. Hence, we have to cover $\mathcal{D}_\zeta$ by smaller basic wide open spaces. As a result, in $\mathcal{D}_\zeta$, we find an affinoid $\mathbf{P}^0_\zeta$, whose reduction is defined by $z_{f+1}^2 = w_1(w_1^{q-1} - 1)^2$. This affine curve has $q-1$ singular points at $w_1 \in k^\times$. Then, by analyzing the tubular neighborhoods of these singular points, we find an affinoid $\mathbf{X}_{\zeta,\zeta'} \subset \mathbf{P}^0_\zeta$ for each $\zeta' \in k^\times$, whose reduction is defined by $z^2 + z = w^3$.

4.4.1 q : odd

We assume that q is odd. For each $\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})$, we define an affinoid $\mathbf{X}_\zeta \subset \mathcal{D}_\zeta$ and compute its reduction $\overline{\mathbf{X}}_\zeta$.

For $\iota \in \mu_2(k^{\mathrm{ac}})$, we choose an element $c'_{1,\iota} \in \mathcal{O}^\times_{K^{\mathrm{ac}}}$ such that $\bar{c}'_{1,\iota} = -2\iota$ and $c'^{2q}_{1,\iota} = 4(1 - \gamma_4 c'_{1,\iota})$. We take $\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})$. We put $c_{1,\zeta} = c'_{1,\zeta^{q^2-1}}$, and define $c_{2,\zeta} \in \mathcal{O}^\times_{K^{\mathrm{ac}}}$ by $c_{2,\zeta}^{q^2-1} = -2c_{1,\zeta}^{-q}$ and $\bar{c}_{2,\zeta} = \zeta$. We put $a_\zeta = \omega_4^{q-1} c_{2,\zeta}^{q+1}$ and $b_\zeta = -2\zeta^{q^2-1}\omega_3^{(q-1)/2} c_{1,\zeta}^{-q} c_{2,\zeta}^{(q+3)/2}$. Note that we have $v(a_\zeta) = 1/(2q^4)$ and $v(b_\zeta) = 1/(4q^3)$.

For an element $\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})$, we define an affinoid $\mathbf{X}_\zeta$ by $v(x_3 - c_{2,\zeta}) \geq 1/(4q^3)$. We change variables as $Z = a_\zeta z + c_{1,\zeta}$, $x_3 = b_\zeta w + c_{2,\zeta}$. Then, we acquire $a_\zeta^q(z^q - z - w^2) \equiv 0 \pmod{\epsilon_3+}$ by (4.27). Dividing this by a_ζ^q, we have $z^q - z = w^2 \pmod{0+}$. Hence, the reduction of $\mathbf{X}_\zeta$ is defined by $z^q - z = w^2$.

Proposition 4.7. *For each* $\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})$, *the reduction* $\overline{\mathbf{X}}_\zeta$ *is defined by* $z^q - z = w^2$ *and the complement* $\mathcal{D}_\zeta \setminus \mathbf{X}_\zeta$ *is an open annulus.*

Proof. We have already proved the first assertion. We prove the second assertion. We change variables as $Z = z' + c_{1,\zeta}$ and $x_3 = w' + c_{2,\zeta}$ with $0 < v(w') < 1/(4q^3)$. Substituting them to (4.27), we obtain $z'^q \equiv w'^2 \pmod{2v(w')+}$. Note that we have $0 < v(z') < 1/(2q^4)$. By setting $w' = z'' z'^{(q-1)/2}$, we acquire $z''^2 \equiv z' \pmod{v(z')+}$. Hence, we can see that z' is written as a function of z''. Then w' is also written as a function of z''. Therefore, $(\mathcal{D}_\zeta \setminus \mathbf{X}_\zeta)(\mathbf{C})$ is identified with $\{z'' \in \mathbf{C} \mid 0 < v(z'') < 1/(4q^4)\}$. ☐

4.4.2 q : even

We assume that q is even. We put $Z_1 = x_3^{q^2-1}$. Then, the congruence (4.27) has the following form;

$$Z^q + Z_1(1 - \gamma_4 Z) + Z_1^{-1} \equiv -\gamma_3^{q^2-q-1} Z_1^{-q(q+1)} \pmod{\epsilon_3+}. \tag{4.28}$$

1. Projective lines For each $\zeta \in k_2^\times$, we define subaffinoid $\mathbf{P}^0_\zeta \subset \mathcal{D}_\zeta$ by $v(Z) \geq 1/(4q^4)$. We change variables as $Z = \varpi^{1/(4q^4)} w_1$ and $Z_1 = 1 + \varpi^{1/(8q^3)} z_1$. Substituting these to (4.28) and dividing it by $\varpi^{1/(4q^3)}$, we acquire

$$(z_1 + w_1^{\frac{q}{2}})^2 + \varpi^{\frac{1}{8q^3}} z_1^3 + \varpi^{\frac{1}{4q^3}} z_1^4 + \varpi^{\frac{q-1}{4q^4}} w_1 \equiv \varpi^{\frac{2q-3}{4q^3}} \pmod{\frac{1}{4q^3}+}. \tag{4.29}$$

We can check that $v(z_1) \geq 0$. We set $q = 2^f$ and put $l_i = (2^i - 1)q/2^i$ and $m_i = 1/(2^{i+2}q^3)$ for $1 \leq i \leq f+1$. Furthermore, we define parameters z_i for $2 \leq i \leq f+1$ by

$$z_i + w_1^{l_i} = \varpi^{m_{i+1}} z_{i+1} \ (1 \leq i \leq f). \tag{4.30}$$

Lemma 4.8. *We assume that $v(Z) \geq 1/(4q^4)$. Then we have*

$$z_{f+1}^2 + w_1^{2q-1} + w_1 + \varpi^{\frac{1}{8q^4}} z_{f+1} w_1^q \equiv \varpi^{\frac{2q^2-4q+1}{4q^4}} \quad (\mathrm{mod}\ \frac{1}{4q^4}+). \tag{4.31}$$

Proof. For $1 \leq i \leq f+1$, we put $n_i = (q - 2^{i-1})/(2^{i+1}q^4)$. We prove

$$(z_i + w_1^{l_i})^2 + \varpi^{m_i} z_i w_1^q + \varpi^{n_i} w_1 \equiv \varpi^{\frac{2^i q - 2^{i+1} + 1}{2^{i+1} q^3}} \quad (\mathrm{mod}\ \frac{1}{2^{i+1}q^3}+) \tag{4.32}$$

for $2 \leq i \leq f+1$ by an induction on i. Eliminating z_1 from (4.29) by (4.30) and dividing it by $\varpi^{1/(8q^4)}$, we obtain

$$(z_2 + w_1^{\frac{3q}{4}})^2 + \varpi^{\frac{1}{16q^3}} z_2 w_1^q + \varpi^{\frac{q-2}{8q^4}} w_1 + \varpi^{\frac{1}{8q^3}} w_1^{\frac{q}{2}} (z_2 + w_1^{\frac{3q}{4}})^2 \equiv \varpi^{\frac{4q-7}{8q^3}} \quad (\mathrm{mod}\ \frac{1}{8q^3}+).$$

This shows $v(z_2 + w_1^{\frac{3q}{4}}) > 1/(32q^3)$. Hence we have (4.32) for $i = 2$. Assuming (4.32) for i. Eliminating z_i from (4.32) by (4.30) and dividing it by ϖ^{m_i}, we obtain (4.32) for $i+1$. Hence, we have (4.32) for $f+1$, which is equivalent to (4.31). □

Proposition 4.9. *For each $\zeta \in k_2^\times$, the reduction $\overline{\mathbf{P}}_\zeta^0$ is an affine curve defined by $z_{f+1}^2 = w_1(w_1^{q-1} - 1)^2$, which has genus 0 and singularities at $w_1 \in k^\times$, and the complement $\mathcal{D}_\zeta \setminus \mathbf{P}_\zeta^0$ is an open annulus.*

Proof. The claim on $\overline{\mathbf{P}}_\zeta^0$ follows from the congruence (4.31) modulo 0+. We prove the last assertion. We change a variable as $Z_1 = 1 + z_1'$ with $0 < v(z_1') < 1/(8q^3)$. Similarly as (4.30), we introduce parameters $\{z_i'\}_{2 \leq i \leq f+1}$ by $z_i' + Z^{l_i} = z_{i+1}'$ for $1 \leq i \leq f$. Then, by similar computations to those in the proof of Lemma 4.8, we obtain $z_{f+1}'^2 \equiv Z^{2q-1} \pmod{2v(z_{f+1}')+}$. By setting $z_{f+2}' = Z^q/z_{f+1}'$, we obtain $z_{f+2}'^2 \equiv Z \pmod{v(Z)+}$. Then we can see that all parameters z_i' for $1 \leq i \leq f+1$ and Z are written as functions of z_{f+2}'. Hence, $(\mathcal{D}_\zeta \setminus \mathbf{P}_\zeta^0)(\mathbf{C})$ is identified with $\{z_{f+2}' \in \mathbf{C} \mid 0 < v(z_{f+2}') < 1/(8q^4)\}$. □

2. Elliptic curves For $\zeta' \in k^\times$, we choose $c_{2,\zeta'} \in \mathcal{O}_{\mathbf{C}}^\times$ such that $\bar{c}_{2,\zeta'} = \zeta'$ and

$$c_{2,\zeta'}^{4(q-1)} + 1 + \varpi^{\frac{1}{4q^4}} c_{2,\zeta'}^{4q-3} = 0,$$

and a square root $c_{2,\zeta'}^{1/2}$ of $c_{2,\zeta'}$. Further, we choose $c_{1,\zeta'}$ such that

$$c_{1,\zeta'}^2 + \varpi^{\frac{1}{8q^4}} c_{2,\zeta'}^q c_{1,\zeta'} + c_{2,\zeta'}(c_{2,\zeta'}^{2(q-1)} + 1) = \varpi^{\frac{2q^2-4q+1}{4q^4}},$$

and $b_{2,\zeta'}$ such that $b_{2,\zeta'}^3 = \varpi^{1/(4q^4)} c_{2,\zeta'}^4$. We put $a_{1,\zeta'} = \varpi^{1/(8q^4)} c_{2,\zeta'}^q$ and $b_{1,\zeta'} = c_{2,\zeta'}^{(2q-3)/2} b_{2,\zeta'}$.

For each $\zeta' \in k^\times$, we define a subspace $\mathcal{D}_{\zeta,\zeta'} \subset \mathbf{P}_\zeta^0$ by $v(w_1 - c_{2,\zeta'}) > 0$. Furthermore, we define $\mathbf{X}_{\zeta,\zeta'} \subset \mathcal{D}_{\zeta,\zeta'}$ by $v(w_1 - c_{2,\zeta'}) \geq 1/(12q^4)$. We put $\mathbf{P}_\zeta = \mathbf{P}_\zeta^0 \setminus \bigcup_{\zeta' \in k^\times} \mathcal{D}_{\zeta,\zeta'}$.

We take $(\zeta, \zeta') \in k_2^\times \times k^\times$ and compute the reduction of $\mathbf{X}_{\zeta,\zeta'}$. In the sequel, we omit the subscript ζ' of $a_{1,\zeta'}$, $b_{1,\zeta'}$, $b_{2,\zeta'}$ $c_{1,\zeta'}$ and $c_{2,\zeta'}$, if there is no confusion. We change variables as $z_{f+1} = a_1 z + b_1 w + c_1$ and $w_1 = b_2 w + c_2$. By substituting these to (4.31), we acquire

$$a_1^2(z^2 + z + w^3) \equiv 0 \quad (\mathrm{mod}\ \frac{1}{4q^4}+) \tag{4.33}$$

by the definition of a_1, b_1, b_2, c_1 and c_2.

Proposition 4.10. *For each $(\zeta, \zeta') \in k_2^\times \times k^\times$, the reduction of $\mathbf{X}_{\zeta,\zeta'}$ is defined by $z^2 + z = w^3$ and the complement $\mathcal{D}_{\zeta,\zeta'} \setminus \mathbf{X}_{\zeta,\zeta'}$ is an open annulus.*

Proof. The first assertion follows from (4.33). We prove the second assertion. We change variables as $z_{f+1} = z' + c_2^{(2q-3)/2} w' + c_1$ and $w_1 = w' + c_2$ with $0 < v(w') < 1/(12q^4)$. Substituting them to (4.31), we acquire $z'^2 \equiv c_2^{2(q-2)} w'^3 \pmod{2v(z')+}$ by the choice of c_2. Note that we have $v(z') = 3v(w')/2 < 1/(8q^4)$. By setting $z'' = z'/(c_2^{q-2} w')$, we obtain $z''^2 \equiv w' \pmod{v(w')+}$. Then we can see that z' and w' are written as functions of z''. Hence, $(\mathcal{D}_{\zeta,\zeta'} \setminus \mathbf{X}_{\zeta,\zeta'})(\mathbf{C})$ is identified with $\{z'' \in \mathbf{C} \mid 0 < v(z'') < 1/(24q^4)\}$. □

4.5 Stable covering of $\mathbf{X}_1(\mathfrak{p}^3)$

In this subsection, we show the existence of the stable covering of $\mathbf{X}_1(\mathfrak{p}^3)$ over some finite extension of the base field $\widehat{K}^{\mathrm{ur}}$. See [CM, Section 2.3] for the notion of semi-stable coverings. A semi-stable coverings is called stable, if the corresponding semi-stable model is stable.

Proposition 4.11. *There exists a stable covering of* $\mathbf{X}_1(\mathfrak{p}^3)$ *over a finite extension of the base field.*

Proof. First, we show that, after taking a finite extension of base field, $\mathbf{X}_1(\mathfrak{p}^3)$ is a wide open space. By [St2, Theorem 2.3.1 (i)], $\mathbf{X}_1(\mathfrak{p}^3)$ is the Raynaud generic fiber of the formal completion of an affine scheme over $\mathcal{O}_{\widehat{K}^{\mathrm{ur}}}$ at a closed point on a special fiber. Then we can apply [CM, Theorem 2.29] to the formal completion of the affine scheme along its special fiber, after shrinking the affine scheme. Hence, $\mathbf{X}_1(\mathfrak{p}^3)$ is a wide open space over some extension.

By [CM, Theorem 2.18], a wide open space can be embedded to a proper algebraic curve so that its complement is a disjoint union of closed disks. Therefore, $\mathbf{X}_1(\mathfrak{p}^3)$ has a semi-stable covering over some finite extension by [CM, Theorem 2.40]. Then a simple modification gives a stable covering. ☐

In the following, we construct a candidate of a semi-stable covering of $\mathbf{X}_1(\mathfrak{p}^3)$ over some finite extension. We put

$$\mathbf{V}_1 = \mathbf{W}_{1,1'}^{+} \cup \bigcup_{2 \le i \le 6} \mathbf{W}_{i,1'}, \ \mathbf{V}_2 = \mathbf{W}_{1,1'}^{-} \cup \bigcup_{2 \le i \le 4} \mathbf{W}_{1,i'}, \ \mathbf{U} = \mathbf{W}_{1,1'} \backslash \bigcup_{\zeta \in \mathcal{S}_1} \mathbf{X}_{\zeta}.$$

We note that $\mathbf{V}_1 \supset \mathbf{Y}_{1,2}$, $\mathbf{V}_2 \supset \mathbf{Y}_{2,1}$, $\mathbf{U} \supset \mathbf{Z}_{1,1}$, $\mathbf{V}_1 \cap \mathbf{V}_2 = \emptyset$, $\mathbf{V}_1 \cap \mathbf{U} = \mathbf{W}_{1,1'}^{+}$ and $\mathbf{V}_2 \cap \mathbf{U} = \mathbf{W}_{1,1'}^{-}$.

We consider the case where q is even in this paragraph. For $\zeta \in k_2^{\times}$, we set $\hat{\mathcal{D}}_{\zeta} = \mathcal{D}_{\zeta} \backslash \big(\bigcup_{\zeta' \in k^{\times}} \mathbf{X}_{\zeta,\zeta'} \big)$. Then, $\hat{\mathcal{D}}_{\zeta}$ contains $\mathbf{P}_{\zeta}$ as the underlying affinoid. On the other hand, for $(\zeta, \zeta') \in k_2^{\times} \times k^{\times}$ the space $\mathcal{D}_{\zeta,\zeta'}$ has the underlying affinoid $\mathbf{X}_{\zeta,\zeta'}$.

We put

$$\mathcal{S} = \begin{cases} \mathcal{S}_1 & \text{if } q \text{ is odd,} \\ k_2^{\times} \times k^{\times} & \text{if } q \text{ is even.} \end{cases}$$

Now, we define a covering of $\mathbf{X}_1(\mathfrak{p}^3)$ as

$$\mathcal{C}_1(\mathfrak{p}^3) = \begin{cases} \{\mathbf{V}_1, \mathbf{V}_2, \mathbf{U}, \{\mathcal{D}_{\zeta}\}_{\zeta \in \mathcal{S}_1}\} & \text{if } q \text{ is odd,} \\ \{\mathbf{V}_1, \mathbf{V}_2, \mathbf{U}, \{\hat{\mathcal{D}}_{\zeta}\}_{\zeta \in k_2^{\times}}, \{\mathcal{D}_{\zeta,\zeta'}\}_{(\zeta,\zeta') \in \mathcal{S}} & \text{if } q \text{ is even.} \end{cases}$$

In Subsection 7.2, we will show that $\mathcal{C}_1(\mathfrak{p}^3)$ is a semi-stable covering of $\mathbf{X}_1(\mathfrak{p}^3)$ over some finite extension.

5 Action of the division algebra on the reductions

In this section, we determine the action of of the division algebra $\mathcal{O}_D^{\times}$ on the reductions $\overline{\mathbf{Y}}_{1,2}$, $\overline{\mathbf{Y}}_{2,1}$ $\overline{\mathbf{Z}}_{1,1}$, $\{\overline{\mathbf{P}}_{\zeta}\}_{\zeta \in k_2^{\times}}$ and $\{\overline{\mathbf{X}}_{\zeta}\}_{\zeta \in \mathcal{S}}$ by using the description of $\mathcal{O}_D^{\times}$-action in (1.8). We take $d = d_1 + \varphi d_2 \in \mathcal{O}_D^{\times}$, where $d_1 \in \mathcal{O}_{K_2}^{\times}$ and $d_2 \in \mathcal{O}_{K_2}$. We put $\kappa_1(d) = \bar{d}_1$ and $\kappa_2(d) = -\bar{d}_2 / \bar{d}_1^q$.

Lemma 5.1. *The element d induces the following morphisms;*

$$\overline{\mathbf{Y}}_{1,2} \to \overline{\mathbf{Y}}_{1,2}; \ (x,y) \mapsto (\kappa_1(d)x, \kappa_1(d)^{-q}y), \ \overline{\mathbf{Y}}_{2,1} \to \overline{\mathbf{Y}}_{2,1}; \ (x,y) \mapsto (\kappa_1(d)^{-1}x, \kappa_1(d)^{q}y).$$

Proof. We prove the assertion for $\overline{\mathbf{Y}}_{1,2}$. By (1.5), we have $d^* x_1 \equiv d_1 x_1$, $d^* x_3 \equiv d_1 x_3 \pmod{0+}$. Therefore, the required assertion follows from (4.5). The assertion for $\overline{\mathbf{Y}}_{1,2}$ is proved similarly. ☐

Now, let the notation be as in subsection 4.3. We put $x_i' = d^* x_i$ for $1 \le i \le 3$, $t' = d^* t$, $Z_0' = d^* Z_0$ and $Z' = d^* Z$. We have $j^{-1}(x_1) \equiv x_1 + d_1^{-q} d_2 \varpi^{\epsilon_1} \tilde{u} x_1$, $j^{-1}(x_2) \equiv x_2 + d_1^{-q} d_2 \varpi^{\epsilon_1} \tilde{u} x_2 \pmod{\epsilon_1 +}$ and $j^{-1}(x_3) \equiv x_3 \pmod{\epsilon_2 +}$ by (1.11). On the other hand, we have $\tilde{d}(x_1) \equiv d_1 x_1$, $\tilde{d}(x_2) \equiv d_1 x_2 + d_2^q \varpi^{\epsilon_1} x_2^q \pmod{\epsilon_1 +}$ and $\tilde{d}^{-1}(x_3) \equiv d_1 x_3 + d_2^q \varpi^{\epsilon_3} x_3^q \pmod{\epsilon_2 +}$ by (1.5). Hence, we obtain

$$x_1' \equiv d_1 x_1 + d_1^{-(q-1)} d_2 \varpi^{\epsilon_1} \tilde{u} x_1, \ x_2' \equiv d_1 x_2 + d_1^{-(q-1)} d_2 \varpi^{\epsilon_1} \tilde{u} x_2 + d_2^q \varpi^{\epsilon_1} x_2^q \pmod{\epsilon_1 +}, \tag{5.1}$$

$$x_3' \equiv d_1 x_3 + d_2^q \varpi^{\epsilon_3} x_3^q \pmod{\epsilon_2 +}. \tag{5.2}$$

By the definition of t and the equation $x_2'/x_1' = -1 + \gamma_2(x_2'^q/t')$, we acquire

$$t' \equiv d_1^q t - d_1^{q-1} d_2^q t^{2-q} \varpi^{\epsilon_2} \pmod{\epsilon_2+} \tag{5.3}$$

using (5.1). We put $G_0 = d_1^{-q} d_2 x_3^{q(q-1)} + d_1^{-1} d_2^q x_3^{-q(q-1)}$. By the definition of Z_0 and the equation $1 + (x_3'^q/t') = \gamma_3 Z_0'$, we obtain

$$Z_0' \equiv Z_0 - \varpi^{\epsilon_3} G_0 \pmod{\epsilon_3+} \tag{5.4}$$

using (5.2) and (5.3). We put $G = G_0 + d_1^{-1} d_2^q (x_2 x_3^{q-2} + x_1^{-1} x_3^q)$. By the definition of Z and the equation $Z_0' + (x_2'/x_3') - (x_3'/x_1') = \gamma_4 (x_2'/x_3') Z'$, we obtain

$$Z' \equiv Z - \frac{x_3}{x_2} \varpi^{\epsilon_4} G \pmod{\epsilon_4+} \tag{5.5}$$

using (5.1), (5.2) and (5.4). We have $G \equiv d_1^{-q} d_2 x_3^{q(q-1)} + d_1^{-1} d_2^q x_3^{(q-1)(q+2)}$ (mod 0+). by $x_1 \equiv -x_3^{q^2}$, $x_2 \equiv x_3^{q^2}$ (mod 0+). We put $\Delta = d_1^{-q} d_2 x_3^{-(q-1)} + d_1^{-1} d_2^q x_3^{q-1}$. Then the congruence (5.5) has the following form;

$$Z' \equiv Z - \varpi^{\epsilon_4} \Delta \pmod{\epsilon_4+}. \tag{5.6}$$

Proposition 5.2. *The element d acts on $\overline{\mathbf{Z}}_{1,1}$ by $(Z, x_3) \mapsto (Z, \kappa_1(d) x_3)$.*

Proof. This follows from (5.2) and (5.6). ☐

Proposition 5.3. *The element d induces the morphism $\overline{\mathbf{P}}_\zeta \to \overline{\mathbf{P}}_{\kappa_1(d)\zeta}$; $w_1 \mapsto w_1$.*

Proof. This follows from (5.6), Proposition 5.2 and $Z = \varpi^{1/(4q^4)} w_1$. ☐

Proposition 5.4. *We take $\zeta \in \mathcal{S}_1$. Further, we take $\zeta' \in k^\times$, if q is even. We set as follows;*

$$\mathcal{S} \ni \eta = \begin{cases} \zeta & \text{if } q \text{ is odd,} \\ (\zeta, \zeta') & \text{if } q \text{ is even,} \end{cases} \quad \mathcal{S} \ni d\eta = \begin{cases} \kappa_1(d)\zeta & \text{if } q \text{ is odd,} \\ (\kappa_1(d)\zeta, \zeta') & \text{if } q \text{ is even,} \end{cases}$$

$$f_d = \begin{cases} \mathrm{Tr}_{k_2/k}(\zeta^{-2q} \kappa_2(d)) & \text{if } q \text{ is odd,} \\ \mathrm{Tr}_{k_2/\mathbb{F}_2}(\zeta^{1-q} \zeta'^{-2} \kappa_2(d)) & \text{if } q \text{ is even.} \end{cases}$$

Then, the element d induces the following morphism

$$\overline{\mathbf{X}}_\eta \to \overline{\mathbf{X}}_{d\eta} \colon \begin{cases} (z, w) \mapsto (\kappa_1(d)^{-(q+1)}(z + f_d), \kappa_1(d)^{-(q+1)/2} w) & \text{if } q \text{ is odd,} \\ (z, w) \mapsto (z + f_d, w) & \text{if } q \text{ is even.} \end{cases}$$

Proof. First, we assume that q is odd. Recall that $Z = a_\zeta z + c_{1,\iota}$ and $x_3 = b_\zeta w + c_{2,\zeta}$. Similarly, we have $Z' = a_{\bar{d}_1\zeta} z' + c_{1,\iota}$ and $x_3 = b_{\bar{d}_1\zeta} w' + c_{2,\bar{d}_1\zeta}$. Then, the claim follows from (5.6).

Next, we assume that q is even. By (5.6) and $d^* x_3 \equiv d_1 x_3$ (mod $(\epsilon_3/2)+$), we acquire

$$d^* z_{f+1} - z_{f+1} \equiv \varpi^{\frac{\epsilon_4}{4}} \sum_{i=1}^{f} (w_1^{q-2^i} \Delta^{2^{i-1}}) \pmod{(\epsilon_4/4)+} \tag{5.7}$$

on the locus where $v(Z) \geq \epsilon_4/2$. By $z_{f+1} = a_{1,\zeta'} z + b_{1,\zeta'} w + c_{1,\zeta'}$ and $w_1 = b_{2,\zeta'} w + c_{2,\zeta'}$, we obtain $d^* z - z \equiv \sum_{i=1}^{f} (c_{2,\zeta'}^{-2^i} \Delta^{2^{i-1}})$ (mod 0+) and $d^* w \equiv w$ (mod $(\epsilon_4/3)+$) on $\mathbf{X}_{\zeta,\zeta'}$ by (5.6) and (5.7). On the other hand, $\sum_{i=1}^{f} (\bar{c}_{2,\zeta'}^{-2^i} \overline{\Delta}^{2^{i-1}}) = f_d$, because $\bar{x_3} = \zeta$ and $\bar{c}_{2,\zeta'} = \zeta'$. Hence, we have proved the claim. ☐

6 Action of the Weil group on the reductions

In this section, we compute the actions of the Weil group on the reductions $\overline{\mathbf{Y}}_{1,2}$, $\overline{\mathbf{Y}}_{2,1}$, $\overline{\mathbf{Z}}_{1,1}$, $\{\overline{\mathbf{P}}_\zeta\}_{\zeta \in k_2^\times}$ and $\{\overline{\mathbf{X}}_\eta\}_{\eta \in \mathcal{S}}$.

Let $\mathbf{X}$ be a reduced affinoid over K^{ur}. For $P \in \mathbf{X}(\mathbf{C})$, the image of P under the natural reduction map $\mathbf{X}(\mathbf{C}) \to \overline{\mathbf{X}}(k^{\mathrm{ac}})$ is denoted by $\overline{P}$. The action of W_K on $\overline{\mathbf{X}}$ is a homomorphism $w_{\mathbf{X}} \colon W_K \to \mathrm{Aut}(\overline{\mathbf{X}})$ characterized by $\overline{\sigma(P)} = w_{\mathbf{X}}(\sigma)(\overline{P})$ for $\sigma \in W_K$ and $P \in \mathbf{X}(\mathbf{C})$. For $\sigma \in W_K$, we define $r_\sigma \in \mathbb{Z}$ so that σ induces q^{-r_σ}-th power map on the residue field of K^{ac}.

Remark 6.1. *In the usual sense, W_K does not act on $\mathbf{X}_1(\mathfrak{p}^3)$, because the action of W_K does not preserve the connected components of $\mathrm{LT}_1(\mathfrak{p}^3)$. Precisely, $w_{\mathbf{X}}$ is an action of $\{(\sigma, \varphi^{-r_\sigma}) \in W_K \times D^\times\}$, which preserves the connected components of $\mathrm{LT}_1(\mathfrak{p}^3)$.*

6.1 Actions of the Weil group on $\overline{\mathbf{Y}}_{1,2}$, $\overline{\mathbf{Y}}_{2,1}$ and $\overline{\mathbf{Z}}_{1,1}$

For $\sigma \in W_K$, we put $\lambda(\sigma) = \overline{\sigma(\varpi^{1/(q^2-1)})/\varpi^{1/(q^2-1)}} \in k_2^{\times}$. We note that λ is not a group homomorphism in general.

Lemma 6.2. *Let $\sigma \in W_K$. Then, the element σ induces the automorphisms*

$$\overline{\mathbf{Y}}_{1,2} \to \overline{\mathbf{Y}}_{1,2};\ (x,y) \mapsto (\lambda(\sigma)^q x^{q^{-r_\sigma}}, \lambda(\sigma)^{-1} y^{q^{-r_\sigma}}),\ \overline{\mathbf{Y}}_{2,1} \to \overline{\mathbf{Y}}_{2,1};\ (x,y) \mapsto (\lambda(\sigma)^{-1} x^{q^{-r_\sigma}}, \lambda(\sigma)^q y^{q^{-r_\sigma}})$$

as schemes over k.

Proof. We prove the claim for $\overline{\mathbf{Y}}_{1,2}$. We set $\sigma(\varpi^{1/(q^3(q^2-1))}) = \xi\varpi^{1/(q^3(q^2-1))}$ with $\xi \in \mu_{q^3(q^2-1)}(K^{\mathrm{ac}})$. Let $P \in \mathbf{Y}_{1,2}(\mathbf{C})$. We have $X_3(\sigma(P)) = \sigma(X_3(P))$. By applying σ to $X_3(P) = \varpi^{1/(q^3(q^2-1))}x_3(P)$, we obtain $x_3(\sigma(P)) = \xi\sigma(x_3(P)) \equiv \xi x_3(P)^{q^{-r_\sigma}} \pmod{0+}$. In the same way, we have $x_1(\sigma(P)) \equiv \xi^{q^4} x_1(P)^{q^{-r_\sigma}} \pmod{0+}$. Therefore, we acquire $x^\sigma = \bar{\xi} x^{q^{-r_\sigma}}$ and $y^\sigma = \bar{\xi}^{-q} y^{q^{-r_\sigma}}$ by (4.5). Hence, the claim follows from $\bar{\xi} = \lambda(\sigma)^q$. We can prove the claim for $\overline{\mathbf{Y}}_{2,1}$ similarly. $\square$

For $\sigma \in W_K$, we put $\xi_\sigma = \sigma(\omega_3)/\omega_3 \in \mu_{2q^3(q-1)}(K^{\mathrm{ac}})$.

Lemma 6.3. *Let $\sigma \in W_K$. Then, σ acts on $\overline{\mathbf{Z}}_{1,1}$ by $(Z, x_3) \mapsto (Z^{q^{-r_\sigma}}, \bar{\xi}_\sigma x_3^{q^{-r_\sigma}})$.*

Proof. We use the notation in Paragraph 4.3. Let $P \in \mathbf{Z}_{1,1}(\mathbf{C})$. Since we set $X_1 = \omega_1^{2q-1}x_1$, $X_2 = \omega_1 x_2$ and $X_3 = \omega_3 x_3$, we have $x_1(\sigma(P)) = \xi_\sigma^{q^2(2q-1)}\sigma(x_1(P))$, $x_2(\sigma(P)) = \xi_\sigma^{q^2}\sigma(x_2(P))$ and $x_3(\sigma(P)) = \xi_\sigma\sigma(x_3(P))$. Hence, we obtain $x_2(\sigma(P))/x_1(\sigma(P)) = \xi_\sigma^{-2q^2(q-1)}\sigma(x_2(P)/x_1(P)) \equiv \sigma(x_2(P)/x_1(P)) \pmod{\epsilon_1+}$. Since we set $x_2/x_1 = -1 + \gamma_2(x_2^q/t)$, we acquire $t(\sigma(P)) \equiv \xi_\sigma^{q^3}\sigma(t(P)) \pmod{\epsilon_2+}$. Therefore, we obtain $(x_3(\sigma(P)))^q/t(\sigma(P)) = \xi_\sigma^{-q(q^2-1)}\sigma(x_3(P)^q/t(P)) \equiv \sigma(x_3(P)^q/t(P)) \pmod{\epsilon_2+}$. Since we set $1 + (x_3^q/t) = \gamma_3 Z_0$, we obtain $Z_0(\sigma(P)) \equiv \sigma(Z_0(P)) \pmod{\epsilon_3+}$. Therefore we acquire

$$Z(\sigma(P)) \equiv \sigma(Z(P)) \pmod{\epsilon_4+} \tag{6.1}$$

by $Z_0 + (x_2/x_3) - (x_3/x_1) = \gamma_4(x_2/x_3)Z$.

The assertion follows from $x_3(\sigma(P)) = \xi_\sigma\sigma(x_3(P)) \equiv \xi_\sigma x_3(P)^{q^{-r_\sigma}} \pmod{0+}$ and (6.1). $\square$

6.2 Action of the Weil group on $\overline{\mathbf{X}}_\eta$

In this subsection, let $\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})$. Until Lemma 6.8, let $\sigma \in W_K$.

6.2.1 q : odd

We assume that q is odd. We use the notation in Paragraph 4.4.1. By (6.1) and $x_3(\sigma(P)) = \xi_\sigma\sigma(x_3(P))$, we have

$$a_{\bar{\xi}_\sigma\zeta^{q^{-r_\sigma}}} z(\sigma(P)) + c_{1,\bar{\xi}_\sigma\zeta^{q^{-r_\sigma}}} = Z(\sigma(P)) \equiv \sigma(Z(P)) = \sigma(a_\zeta)\sigma(z(P)) + \sigma(c_{1,\zeta}) \pmod{\epsilon_4+}, \tag{6.2}$$

$$b_{\bar{\xi}_\sigma\zeta^{q^{-r_\sigma}}} w(\sigma(P)) + c_{2,\bar{\xi}_\sigma\zeta^{q^{-r_\sigma}}} = x_3(\sigma(P)) = \xi_\sigma\sigma(x_3(P)) = \xi_\sigma\sigma(b_\zeta)\sigma(w(P)) + \xi_\sigma\sigma(c_{2,\zeta}) \tag{6.3}$$

for $P \in \mathbf{X}_\zeta(\mathbf{C})$. Note that $c_{1,\bar{\xi}_\sigma\zeta^{q^{-r_\sigma}}} = c_{1,\zeta}$ and $c_{2,\bar{\xi}_\sigma\zeta^{q^{-r_\sigma}}} = \xi_\sigma^{q^4}\zeta^{q^{-r_\sigma}-1}c_{2,\zeta}$. We have $v(\sigma(c_{1,\zeta}) - c_{1,\zeta}) \geq \epsilon_4$ by (6.2). We put

$$a_{\sigma,\zeta} = \frac{\sigma(a_\zeta)}{\zeta^{r_\sigma(q^2-1)}\xi_\sigma^{q+1}a_\zeta},\ b_{\sigma,\zeta} = \left(\frac{\sigma(c_{1,\zeta}) - c_{1,\zeta}}{\zeta^{r_\sigma(q^2-1)}\xi_\sigma^{q+1}a_\zeta}\right),\ c_{\sigma,\zeta} = \frac{\sigma(b_\zeta)}{\xi_\sigma^{\frac{q+1}{2}}b_\zeta}.$$

Then we have $a_{\sigma,\zeta}, b_{\sigma,\zeta}, c_{\sigma,\zeta} \in \mathcal{O}_{K^{\mathrm{ac}}}$. In the sequel, we omit the subscript ζ of $a_{\sigma,\zeta}$, $b_{\sigma,\zeta}$ and $c_{\sigma,\zeta}$.

Proposition 6.4. *We have $\bar{a}_\sigma \in k^\times$, $\bar{b}_\sigma \in k$ and $\bar{a}_\sigma = \bar{c}_\sigma^2$. Further, σ induces the morphism*

$$\overline{\mathbf{X}}_\zeta \to \overline{\mathbf{X}}_{\bar{\xi}_\sigma\zeta};\ (z,w) \mapsto (\bar{a}_\sigma z^{q^{-r_\sigma}} + \bar{b}_\sigma, \bar{c}_\sigma w^{q^{-r_\sigma}}).$$

Proof. We have $v(\xi_\sigma\sigma(c_{2,\zeta}) - \xi_\sigma^{q^4}\zeta^{q^{-r_\sigma}-1}c_{2,\zeta}) \geq \epsilon_3$ by $v(\sigma(c_{1,\zeta}) - c_{1,\zeta}) \geq \epsilon_4$. Hence we have the last assertion by (6.2) and (6.3). By the definition of a_ζ, b_ζ and $c_{1,\zeta}$, we can check that $\bar{a}_\sigma^{q-1} = 1$, $\bar{b}_\sigma^q = \bar{b}_\sigma$ and $\bar{a}_\sigma = \bar{c}_\sigma^2$ using $c_{1,\zeta}^q \equiv -\iota(2 - \gamma_4 c_{1,\zeta}) \pmod{(q-1)/q^4}$. $\square$

We put $L = K(\varpi^{1/2})$ and $L_2 = K_2(\varpi^{1/2})$ in K^{ac}. Let LT_{L_2} be the universal formal $\mathcal{O}_L$-module over L^{ur} of dimension 1 and height 1. We have $[\varpi^{1/2}]_{\mathrm{LT}_{L_2}}(X) = \varpi^{1/2}X - X^{q^2}$. We put $\varpi_{1,L_2} = \varpi^{1/(2(q^2-1))}$ and take $\varpi_{2,L_2} \in \mathcal{O}_{K^{\mathrm{ac}}}$ such that $[\varpi^{1/2}]_{\mathrm{LT}_{L_2}}(\varpi_{2,L_2}) = \varpi_{1,L_2}$. Let $\mathrm{Art}_{L_2} \colon L_2^{\times} \xrightarrow{\sim} W_{L_2}^{\mathrm{ab}}$ be the Artin reciprocity map normalized so that the image by Art_{L_2} of a uniformizer is a lift of the geometric Frobenius. We consider the following homomorphism;

$$I_{L_2} \to k_2^{\times} \times k_2;\ \sigma \mapsto \bigl(\lambda_1(\sigma), \lambda_2(\sigma)\bigr) = \left(\overline{\frac{\sigma(\varpi_{1,L_2})}{\varpi_{1,L_2}}}, \overline{\left(\frac{\varpi_{1,L_2}\sigma(\varpi_{2,L_2}) - \sigma(\varpi_{1,L_2})\varpi_{2,L_2}}{\sigma(\varpi_{1,L_2})\varpi_{1,L_2}}\right)}\right).$$

This map is equal to the composite $I_{L_2} \to \mathcal{O}_{L_2}^{\times} \to (\mathcal{O}_{L_2}/\varpi)^{\times} \simeq k_2^{\times} \times k_2$, where the first homomorphism is induced from the inverse of Art_{L_2}. Then, we rewrite Proposition 6.4 as follows:

Corollary 6.5. *Let $\sigma \in I_L$. We put $g_0 = (2/\zeta^{q+1})(\lambda_2(\sigma)^q + \zeta^{q^2-1}\lambda_2(\sigma)) \in k$. Then, σ induces the morphism*

$$\overline{\mathbf{X}}_{\zeta} \to \overline{\mathbf{X}}_{\lambda_1(\sigma)^{q+1}\zeta};\ (z, w) \mapsto (\lambda_1(\sigma)^{-2(q+1)}(z + g_0), \lambda_1(\sigma)^{-(q+1)}w).$$

Proof. We can check that $\bar{a}_\sigma = \lambda_1(\sigma)^{-2(q+1)}$ and $\bar{c}_\sigma = \lambda_1(\sigma)^{-(q+1)}$ easily. We prove $\bar{b}_\sigma = \lambda_1(\sigma)^{-2(q+1)}g_0$. We simply write ϖ_i for ϖ_{i,L_2}. We put $\iota = \zeta^{q^2-1}$ and $C = \varpi_1^{(q^2-1)/q}\{(\varpi_2/\varpi_1)^q + \iota(\varpi_2/\varpi_1)\}$. Then, we have $C^q - \iota\gamma_1 C \equiv -1 \pmod{(1/2)+}$ by $\varpi_2^{q^2} - \varpi^{1/2}\varpi_2 = -\varpi_1$. We can easily check the equality $\sigma(C) - C \equiv \varpi^{\epsilon_1}(\lambda_2(\sigma)^q + \iota\lambda_2(\sigma)) \pmod{\epsilon_1+}$. On the other hand, we can check $c_{1,\zeta}^q \equiv -\iota(2 - \gamma_4 c_{1,\zeta}) \pmod{((q-1)/(2q^4))+}$ by the definition of $c_{1,\zeta}$. Therefore, the elements C and $c_{1,\zeta}^{q^3}/(2\iota)$ satisfy $x^q - \iota\gamma_1 x \equiv -1 \pmod{(1/2)+}$. Hence, we obtain $C \equiv c_{1,\zeta}^{q^3}/(2\iota) \pmod{\epsilon_1+}$. This implies $(\sigma(c_{1,\zeta}) - c_{1,\zeta})^{q^3} \equiv 2\iota(\sigma(C) - C) \pmod{\epsilon_1+}$. Therefore, we obtain $\bar{b}_\sigma \equiv \bar{b}_\sigma^{q^3} \equiv \lambda_1(\sigma)^{-2(q+1)}g_0 \pmod{0+}$ by $\xi_\sigma = \lambda_1(\sigma)^{q+1} \pmod{0+}$. □

6.2.2 q : even

We assume that q is even. We use the notation in Paragraph 4.4.2. For $P \in \mathbf{P}^0(\mathbf{C})$, we have

$$w_1(\sigma(P)) \equiv \sigma(w_1(P)) \pmod{\frac{1}{4q^4}+} \tag{6.4}$$

by (6.1). We can see that

$$z_{f+1}(\sigma(P)) \equiv \sigma(z_{f+1}(P)) \pmod{\frac{1}{8q^4}+} \tag{6.5}$$

using (4.31) and (6.4).

Lemma 6.6. *The element σ induces the morphism $\overline{\mathbf{P}}_{\zeta} \to \overline{\mathbf{P}}_{\bar{\xi}_\sigma\zeta};\ w_1 \mapsto w_1^{q^{-r_\sigma}}$.*

Proof. This follows from Lemma 6.3 and (6.4). □

We take $\zeta' \in k^{\times}$. By (6.4) and (6.5), we have

$$a_{1,\zeta'}z(\sigma(P)) + b_{1,\zeta'}w(\sigma(P)) + c_{1,\zeta'} \equiv a_{1,\zeta'}\sigma(z(P)) + \sigma(b_{1,\zeta'})\sigma(w(P)) + \sigma(c_{1,\zeta'}) \pmod{\frac{1}{8q^4}+}, \tag{6.6}$$

$$b_{2,\zeta'}w(\sigma(P)) + c_{2,\zeta'} \equiv \sigma(b_{2,\zeta'})\sigma(w(P)) + \sigma(c_{2,\zeta'}) \pmod{\frac{1}{4q^3}+} \tag{6.7}$$

using $\sigma(a_{1,\zeta'}) \equiv a_{1,\zeta'} \pmod{1/(8q^4)+}$. We put

$$a_{\sigma,\zeta'} = \frac{\sigma(b_{2,\zeta'})}{b_{2,\zeta'}},\ b_{\sigma,\zeta'} = \frac{\sigma(b_{1,\zeta'})b_{2,\zeta'} - b_{1,\zeta'}\sigma(b_{2,\zeta'})}{a_{1,\zeta'}b_{2,\zeta'}},$$

$$b'_{\sigma,\zeta'} = \frac{\sigma(c_{2,\zeta'}) - c_{2,\zeta'}}{b_{2,\zeta'}},\ c_{\sigma,\zeta'} = \frac{\sigma(c_{1,\zeta'}) - c_{1,\zeta'} - b_{1,\zeta'}b_{2,\zeta'}^{-1}(\sigma(c_{2,\zeta'}) - c_{2,\zeta'})}{a_{1,\zeta'}}.$$

In the sequel, we omit the subscript ζ' of $a_{\sigma,\zeta'}$, $b_{\sigma,\zeta'}$, $b'_{\sigma,\zeta'}$ and $c_{\sigma,\zeta'}$. We note that $v(a_\sigma) = 0$. We have $v(b'_\sigma) \geq 0$ by (6.7). This implies $v(b_\sigma) \geq 0$. By (6.6) and (6.7), we obtain $v(c_\sigma) \geq 0$ using $v(b_\sigma) \geq 0$.

Proposition 6.7. *The element σ induces the morphism*

$$\overline{\mathbf{X}}_{\zeta,\zeta'} \to \overline{\mathbf{X}}_{\bar{\xi}_\sigma \zeta^{q^{-r_\sigma}},\zeta'};\ (z,w) \mapsto (z^{q^{-r_\sigma}} + \bar{b}_\sigma w^{q^{-r_\sigma}} + \bar{c}_\sigma, \bar{a}_\sigma w^{q^{-r_\sigma}} + \bar{b}'_\sigma).$$

Proof. This follows from (6.6), (6.7). ☐

In the following, we simplify the description of $\bar{a}_\sigma$, $\bar{b}_\sigma$, $\bar{b}'_\sigma$ and $\bar{c}_\sigma$. Let $\tilde{\zeta}' \in \mu_{q-1}(K)$ be the lift of ζ'. We put $h_{\zeta'}(x) = x^4 - \varpi^{1/4}\tilde{\zeta}'^4 x - \tilde{\zeta}'^4$.

Lemma 6.8. *There is a root δ_1 of $h_{\zeta'}(x) = 0$ such that $\delta_1 \equiv c_{2,\zeta'}^{q^4}$ (mod 1/4).*

Proof. We put $h(x) = x^{4(q-1)} + 1 + \varpi^{1/4} x^{4q-3}$. By the definition of $c_{2,\zeta'}$, we have $h(c_{2,\zeta'}^{q^4}) \equiv 0 \pmod 1$. Hence, we have a root c'_2 of h such that $c'_2 \equiv c_{2,\zeta'}^{q^4}$ (mod 3/4) by Newton's method. We can check that $c'_2 \equiv \tilde{\zeta}'$ (mod 1/16). We define a parameter s with $v(s) \geq 1/16$ by $x = \tilde{\zeta}' + s$. Then we have

$$(1 - \tilde{\zeta}'^{-4} s^4) h(\tilde{\zeta}' + s) \equiv \tilde{\zeta}'^{-4} s^4 + \varpi^{\frac{1}{4}}(s + \tilde{\zeta}') \equiv \tilde{\zeta}'^{-4} h_{\zeta'}(x) \pmod{1/2}.$$

This implies $h_{\zeta'}(c'_2) \equiv 0$ (mod 1/2). Therefore, we have a root δ_1 of $h_{\zeta'}(x) = 0$ such that $\delta_1 \equiv c_{2,\zeta'}$ (mod 1/4) by Newton's method. ☐

By the definition of $b_{2,\zeta'}$, we have $b_{2,\zeta'}^{3q^4} \varpi^{-1/4} \equiv \tilde{\zeta}'^4$ (mod 0+). Let ζ'' be the element of $\mu_{3(q-1)}(K^{\mathrm{ur}})$ satisfying $\zeta'' \equiv b_{2,\zeta'}^{q^4} \varpi^{-1/12}$ (mod 0+). Note that $\zeta''^3 = \tilde{\zeta}'^4$. We take δ_1 as in Lemma 6.8 and put $\delta = \delta_1/(\zeta'' \varpi^{1/12})$. Then we have $\delta^4 - \delta = 1/(\zeta'' \varpi^{1/3})$. Note that $v(\delta) = -1/12$. We take $\zeta_3 \in \mu_3(K^{\mathrm{ur}})$ such that $\zeta_3 \neq 1$, and put $h_{\delta_1}(x) = x^2 - (1 + 2\zeta_3)\varpi^{1/4}\delta_1^{2q} x - \varpi^{1/4}\delta_1^{4q-1}(1 + 2\varpi^{1/4}\delta_1)$.

Lemma 6.9. *There is a root θ_1 of $h_{\delta_1}(x) = 0$ such that $\theta_1 \equiv c_{1,\zeta'}^{2q^4}$ (mod 1/4).*

Proof. By the definition of $c_{1,\zeta'}$ and $c_{2,\zeta'}$, we have $h_{\delta_1}(c_{1,\zeta'}^{2q^4}) \equiv 0$ (mod 1/2). Hence, we can show the claim using Newton's method. ☐

We take θ_1 as in Lemma 6.9 and put $\theta = \theta_1/(\varpi^{1/4}\delta_1^{2q}) - \zeta_3$. Then we have $\theta^2 - \theta = \delta^3$. Note that $v(\theta) = -1/8$. Let $\sigma \in W_K$ in this paragraph. We put $\zeta_{3,\sigma} = \sigma(\zeta'' \varpi^{1/3})/(\zeta'' \varpi^{1/3})$. We take $\nu_\sigma \in \mu_3(K^{\mathrm{ur}}) \cup \{0\}$ such that $\sigma(\delta) \equiv \zeta_{3,\sigma}^{-1}(\delta + \nu_\sigma)$ (mod 5/6). Then we have

$$\bigl(\sigma(\theta) - \theta + \nu_\sigma^2 \delta\bigr)^2 \equiv \sigma(\theta) - \theta + \nu_\sigma^2 \delta + \nu_\sigma^3,\ \bigl(\sigma(\theta) - \theta + \nu_\sigma^2 \delta + \nu_\sigma^3\bigr)^2 \equiv \sigma(\theta) - \theta + \nu_\sigma^2 \delta \pmod{0+}. \tag{6.8}$$

By these equations, we can take $\mu_\sigma \in \mu_3(K^{\mathrm{ur}}) \cup \{0\}$ such that $\mu_\sigma \equiv \sigma(\theta) - \theta + \nu_\sigma^2 \delta + \nu_\sigma^3 + \sigma(\zeta_3) - \zeta_3$ (mod 0+). Then we have $\mu_\sigma^2 + \mu_\sigma \equiv \nu_\sigma^3$ (mod 1) by (6.8) and $\nu_\sigma, \mu_\sigma \in \mu_3(K^{\mathrm{ur}}) \cup \{0\}$.

Lemma 6.10. *1. Let $\sigma \in W_K$. Then $a_\sigma \equiv \zeta_{3,\sigma}$, $b_\sigma \equiv \zeta_{3,\sigma}\nu_\sigma^2$, $b'_\sigma \equiv \nu_\sigma$, $c_\sigma \equiv \mu_\sigma$ (mod 0+).*
2. Let $\sigma \in W_K$. Then $\bar{a}_\sigma \in \mathbb{F}_4^\times$ and $\bar{b}_\sigma, \bar{b}'_\sigma, \bar{c}_\sigma \in \mathbb{F}_4$. Further, $\bar{a}_\sigma \bar{b}_\sigma^2 = \bar{b}'_\sigma$ and $\bar{b}_\sigma^3 = \bar{c}_\sigma^2 + \bar{c}_\sigma$ hold.

Proof. By the definition of $b_{2,\zeta'}$, we have $a_\sigma^{4q^4} \equiv \sigma(\zeta''^4 \varpi^{1/3})/(\zeta''^4 \varpi^{1/3})$ (mod 0+). Hence we have $\bar{a}_\sigma^{4q^4} = \bar{\zeta}_{3,\sigma} \in \mathbb{F}_4^\times$. This implies $\bar{a}_\sigma = \bar{\zeta}_{3,\sigma} \in \mathbb{F}_4^\times$.

By the definition of $a_{1,\zeta'}$ and $b_{1,\zeta'}$, we have

$$\begin{aligned} b_\sigma^{2q^4} &\equiv \frac{\sigma(b_{2,\zeta'}^{2q^4})\bigl(\sigma(c_{2,\zeta'}^{q^4(2q-3)}) - c_{2,\zeta'}^{q^4(2q-3)}\bigr)}{\varpi^{\frac{1}{4}} c_{2,\zeta'}^{2q^5}} \equiv \frac{a_\sigma^{2q^4} b_{2,\zeta'}^{2q^4}\bigl(\sigma(\delta_1^{2q-3}) - \delta_1^{2q-3}\bigr)}{\varpi^{\frac{1}{4}}\delta_1^{2q}} \\ &\equiv \frac{\zeta_{3,\sigma}^2 \zeta''^2\bigl(\sigma(\delta_1) - \delta_1\bigr)}{\varpi^{\frac{1}{12}}\delta_1^4} \equiv \zeta_{3,\sigma}^2 \left(\frac{\sigma(\zeta'' \varpi^{\frac{1}{12}})}{\zeta'' \varpi^{\frac{1}{12}}}\sigma(\delta) - \delta\right) \equiv \zeta_{3,\sigma}^2 \nu_\sigma \pmod{0+}, \end{aligned}$$

where we use Lemma 6.8 in the second congruence, $b_{2,\zeta'}^{q^4}/\varpi^{1/12} \equiv \zeta''$ (mod 0+) in the third congruence, $\delta_1^4 = \tilde{\zeta}'^4$ (mod 1/4) and $\zeta''^3 = \tilde{\zeta}'^4$ in the fourth congruence and $\sigma(\zeta'' \varpi^{1/12})/(\zeta'' \varpi^{1/12}) \equiv \zeta_{3,\sigma}$ (mod 0+) in the last congruence. Hence, we obtain $\bar{b}_\sigma = \bar{\zeta}_{3,\sigma}\bar{\nu}_\sigma^2 \in \mathbb{F}_4$.

By Lemma 6.8 and $b_{2,\zeta'}^{q^4}/\varpi^{1/12} \equiv \zeta''$ (mod 0+), we have

$$b'^{q^4}_\sigma \equiv \frac{\sigma(c_{2,\zeta'}^{q^4}) - c_{2,\zeta'}^{q^4}}{b_{2,\zeta'}^{q^4}} \equiv \frac{\sigma(\delta_1) - \delta_1}{\zeta'' \varpi^{\frac{1}{12}}} = \frac{\sigma(\zeta'' \varpi^{\frac{1}{12}})}{\zeta'' \varpi^{\frac{1}{12}}}\sigma(\delta) - \delta \equiv \nu_\sigma \pmod{0+}.$$

Hence, we obtain $\bar{b}'_\sigma = \bar{\nu}_\sigma \in \mathbb{F}_4$.

By Lemma 6.8, Lemma 6.9 and the definition of $a_{1,\zeta'}$, we have

$$\begin{aligned} c_\sigma^{2q^4} &\equiv \frac{\sigma(\theta_1) - \theta_1 - \delta_1^{2q-3}(\sigma(\delta_1^2) - \delta_1^2)}{\varpi^{\frac{1}{4}}\delta_1^{2q}} \equiv \frac{\delta_1^{-2q}\sigma\bigl(\delta_1^{2q}\varpi^{\frac{1}{4}}(\theta+\zeta_3)\bigr) - \varpi^{\frac{1}{4}}(\theta+\zeta_3) - \delta_1^{-3}(\sigma(\delta_1)-\delta_1)^2}{\varpi^{\frac{1}{4}}} \\ &\equiv \frac{\sigma\bigl(\varpi^{\frac{1}{4}}(\theta+\zeta_3)\bigr) - \varpi^{\frac{1}{4}}(\theta+\zeta_3) - \varpi^{\frac{1}{12}}\delta\bigl(\sigma(\varpi^{\frac{1}{12}}\delta) - \varpi^{\frac{1}{12}}\delta\bigr)^2}{\varpi^{\frac{1}{4}}} \\ &\equiv \sigma(\theta) - \theta + \nu_\sigma^2\delta + \sigma(\zeta_3) - \zeta_3 \pmod{0+}, \end{aligned}$$

where we use $\sigma(\delta_1) \equiv \delta_1 \pmod{1/4}$ in the second congruence, $\delta_1^4 = \tilde{\zeta}'^4 \pmod{1/4}$ in the third congruence. Then we have $\bar{c}_\sigma^{2q^4} \in \mathbb{F}_4$ by (6.8). Hence we have $\bar{c}_\sigma \in \mathbb{F}_4$ and $c_\sigma \equiv \mu_\sigma \pmod{0+}$ again by (6.8).

By the above calculations, we can easily check $\bar{a}_\sigma \bar{b}_\sigma^2 = \bar{b}'_\sigma$ and $\bar{b}_\sigma^3 = \bar{c}_\sigma^2 + \bar{c}_\sigma$. ☐

Lemma 6.11. *The field $K(\zeta_3, \zeta''\varpi^{1/3}, \theta)$ is a Galois extension over K.*

Proof. Let $\sigma \in W_K$. It suffices to show $\sigma(\theta) \in K(\zeta_3, \zeta''\varpi^{1/3}, \theta)$. We put $\theta_\sigma = \theta + \nu_\sigma^2\delta + \nu_\sigma^3 + \mu_\sigma + \sigma(\zeta_3) - \zeta_3$. Then we have $\theta_\sigma^2 - \theta_\sigma \equiv \sigma(\delta)^3 \pmod{2/3}$. Hence, we can find θ' such that $\theta'^2 - \theta' = \sigma(\delta)^3$ and $\theta' \equiv \theta_\sigma \pmod{2/3}$. By the choice of μ_σ, we have $\theta' = \sigma(\theta) \pmod{0+}$. Hence, we obtain $\theta' = \sigma(\theta)$.

We take $\sigma' \in W_K$ such that $\sigma'(\theta) \neq \sigma(\theta)$. We can define $\theta_{\sigma'}$ as above, and have $\sigma'(\theta) \equiv \theta_{\sigma'} \pmod{2/3}$. If $\nu_\sigma = \nu_{\sigma'}$, then we have $\zeta_{3,\sigma}\sigma(\delta) \equiv \zeta_{3,\sigma'}\sigma'(\delta) \pmod{5/6}$, which implies $\zeta_{3,\sigma}\sigma(\delta) = \zeta_{3,\sigma'}\sigma'(\delta)$ because both sides are roots of $x^4 - x - 1/(\zeta''\varpi^{1/3}) = 0$. Hence, if $\sigma(\delta)^3 \neq \sigma'(\delta)^3$, we have $\nu_\sigma \neq \nu_{\sigma'}$, which implies $\sigma(\theta) \equiv \theta_\sigma \not\equiv \theta_{\sigma'} \equiv \sigma'(\theta) \pmod{0+}$. If $\sigma(\delta)^3 = \sigma'(\delta)^3$, we have $\sigma(\theta) \not\equiv \sigma'(\theta) \pmod{0+}$. Therefore we have $v(\sigma(\theta) - \theta_\sigma) > v(\sigma'(\theta) - \theta_\sigma)$. Then, we obtain $\sigma(\theta) \in K(\theta_\sigma) \subset K(\zeta_3, \zeta''\varpi^{1/3}, \theta)$ by Krasner's lemma. ☐

Let E be the elliptic curve over k^{ac} defined by $z^2 + z = w^3$. We put

$$Q = \left\{ g(\alpha,\beta,\gamma) = \begin{pmatrix} \alpha & \beta & \gamma \\ & \alpha^2 & \beta^2 \\ & & \alpha \end{pmatrix} \in GL_3(\mathbb{F}_4) \;\middle|\; \alpha\gamma^2 + \alpha^2\gamma = \beta^3 \right\}.$$

We note that $|Q| = 24$ and Q is isomorphic to $SL_2(\mathbb{F}_3)$ (cf. [Se, 8.5. Exercices 2]). Let $Q \rtimes \mathbb{Z}$ be a semidirect product, where $r \in \mathbb{Z}$ acts on Q by $g(\alpha,\beta,\gamma) \mapsto g(\alpha^{q^r}, \beta^{q^r}, \gamma^{q^r})$. Then $Q \rtimes \mathbb{Z}$ acts faithfully on E as a scheme over k, where $(g(\alpha,\beta,\gamma), r) \in Q \rtimes \mathbb{Z}$ acts on E by $(z,w) \mapsto \bigl(z^{q^{-r}} + \alpha^{-1}\beta w^{q^{-r}} + \alpha^{-1}\gamma, \alpha(w^{q^{-r}} + (\alpha^{-1}\beta)^2)\bigr)$ for k^{ac} valued points.

Proposition 6.12. *The element $\sigma \in W_K$ sends $\overline{\mathbf{X}}_{\zeta,\zeta'}$ to $\overline{\mathbf{X}}_{\bar{\xi}_\sigma \zeta^{q^{-r_\sigma}},\zeta'}$. We identify $\overline{\mathbf{X}}_{\zeta,\zeta'}$ with $\overline{\mathbf{X}}_{\bar{\xi}_\sigma \zeta^{q^{-r_\sigma}},\zeta'}$ by $(z,w) \mapsto (z,w)$. Then the action of W_K gives a homomorphism*

$$\Theta_{\zeta'} \colon W_K \to Q \rtimes \mathbb{Z} \subset \mathrm{Aut}_k(\overline{\mathbf{X}}_{\zeta,\zeta'});\ \sigma \mapsto \bigl(g(\bar{\zeta}_{3,\sigma}, \bar{\zeta}_{3,\sigma}^2\bar{\nu}_\sigma^2, \bar{\zeta}_{3,\sigma}\bar{\mu}_\sigma), r_\sigma\bigr).$$

Proof. This follows from Proposition 6.7 and Lemma 6.10. ☐

Proposition 6.13. *The homomorphism $\Theta_{\zeta'}$ factors through $W(K^{\mathrm{ur}}(\varpi^{1/3},\theta)/K)$ and gives an isomorphism $W(K^{\mathrm{ur}}(\varpi^{1/3},\theta)/K) \simeq Q \rtimes \mathbb{Z}$.*

Proof. By Lemma 6.10.1, the homomorphism $\Theta_{\zeta'}$ factors through $W(K^{\mathrm{ur}}(\varpi^{1/3},\theta)/K)$ and induces an injective homomorphism $W(K^{\mathrm{ur}}(\varpi^{1/3},\theta)/K) \to Q \rtimes \mathbb{Z}$.

To prove the surjectivity, it suffices to show that $\Theta_{\zeta'}$ sends I_K onto Q. Let $g = g(\alpha,\beta,\gamma) \in Q$. We take $\zeta_\alpha \in \mu_3(K^{\mathrm{ur}})$, $\nu_\beta, \mu_\gamma \in \mu_3(K^{\mathrm{ur}}) \cup \{0\}$ such that $\bar{\zeta}_\alpha = \alpha$, $\bar{\nu}_\beta = \alpha^{-1}\beta$ and $\bar{\mu}_\gamma = \alpha^{-1}\gamma$. We put $\delta_g = \zeta_\alpha^{-1}(\delta + \nu_\beta)$ and $\theta_g = \theta + \nu_\beta^2\delta + \nu_\beta^3 + \mu_\gamma$. Then we have $\delta_g^4 - \delta_g \equiv 1/(\zeta_\alpha\zeta''\varpi^{1/3}) \pmod{5/6}$. Hence, we can find δ'_g such that $\delta'^4_g - \delta'_g = 1/(\zeta_\alpha\zeta''\varpi^{1/3})$ and $\delta'_g \equiv \delta_g \pmod{5/6}$. Further, we have $\theta_g^2 - \theta_g \equiv \delta'^3_g \pmod{2/3}$. Hence, we can find θ'_g such that $\theta'^2_g - \theta'_g = \delta'^3_g$ and $\theta'_g \equiv \theta_g \pmod{2/3}$. Then $\varpi^{1/3} \mapsto \zeta_\alpha\varpi^{1/3}$ and $\theta \mapsto \theta'_g$ gives an element of I_K, whose image by $\Theta_{\zeta'}$ is g. ☐

7 Cohomology of $\mathbf{X}_1(\mathfrak{p}^3)$

In this section we show that the covering $\mathcal{C}_1(\mathfrak{p}^3)$ is semi-stable, and study a structure of ℓ-adic cohomology of $\mathbf{X}_1(\mathfrak{p}^3)$. In the sequel, for a projective smooth curve X over k, we simply write $H^1(X, \overline{\mathbb{Q}}_\ell)$ for $H^1(X_{\bar{k}}, \overline{\mathbb{Q}}_\ell)$. For a finite abelian group A, the character group $\mathrm{Hom}_{\mathbb{Z}}(A, \overline{\mathbb{Q}}_\ell^\times)$ is denoted by $A^\vee$.

7.1 Cohomology of reductions

Let X_{DL} be the smooth compactification of an affine curve over k defined by $X^q - X = Y^{q+1}$. The curve X_{DL} is also the smooth compactification of the Deligne-Lusztig curve $x^q y - xy^q = 1$ for $SL_2(\mathbb{F}_q)$. Then, $a \in k$ acts on X_{DL} by $\alpha_a \colon (X, Y) \mapsto (X + a, Y)$. On the other hand, $\zeta \in k_2^\times$ acts on X_{DL} by $\beta_\zeta \colon (X, Y) \mapsto (\zeta^{q+1} X, \zeta Y)$. By these actions, we consider $H^1(X_{\mathrm{DL}}, \overline{\mathbb{Q}}_\ell)$ as a $\overline{\mathbb{Q}}_\ell[k \times k_2^\times]$-module.

Lemma 7.1. *We have an isomorphism*

$$H^1(X_{\mathrm{DL}}, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\psi \in k^\vee \backslash \{1\}} \bigoplus_{\chi \in \mu_{q+1}(k_2)^\vee \backslash \{1\}} \psi \otimes \chi$$

as $\overline{\mathbb{Q}}_\ell[k \times \mu_{q+1}(k_2)]$*-modules.*

Proof. As $\overline{\mathbb{Q}}_\ell[k \times \mu_{q+1}(k_2)]$-modules, we have the short exact sequence

$$0 \to \bigoplus_{\psi \in k^\vee} \psi \to H_c^1(X_{\mathrm{DL}} \backslash X_{\mathrm{DL}}(k), \overline{\mathbb{Q}}_\ell) \to H^1(X_{\mathrm{DL}}, \overline{\mathbb{Q}}_\ell) \to 0. \tag{7.1}$$

Let $\mathcal{L}_\psi$ denote the Artin-Schreier $\overline{\mathbb{Q}}_\ell$-sheaf associated to $\psi \in k^\vee$. Let $\mathcal{K}_\chi$ denote the Kummer $\overline{\mathbb{Q}}_\ell$-sheaf associated to $\chi \in \mu_{q+1}(k_2)^\vee$. Since $X_{\mathrm{DL}} \backslash X_{\mathrm{DL}}(k) \to \mathbb{G}_m$; $(X, Y) \mapsto Y^{q+1}$ is a finite etale Galois covering with a Galois group $k \times \mu_{q+1}(k_2)$, we have the isomorphism

$$H_c^1(X_{\mathrm{DL}} \backslash X_{\mathrm{DL}}(k), \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\psi \in k^\vee} \bigoplus_{\chi \in \mu_{q+1}(k_2)^\vee} H_c^1(\mathbb{G}_m, \mathcal{L}_\psi \otimes \mathcal{K}_\chi) \tag{7.2}$$

as $\overline{\mathbb{Q}}_\ell[k \times \mu_{q+1}(k_2)]$-modules. Note that we have $\dim H_c^1(\mathbb{G}_m, \mathcal{L}_\psi \otimes \mathcal{K}_\chi) = 1$ if $\psi \neq 1$ by the Grothendieck-Ogg-Shafarevich formula (cf. [SGA5, Exposé X Théorème 7.1]). Clearly, if $\chi \neq 1$, we have $H_c^1(\mathbb{G}_m, \mathcal{K}_\chi) = 0$ and $H_c^1(\mathbb{G}_m, \mathcal{L}_\psi) \simeq \psi$. Hence, we acquire the isomorphism

$$\bigoplus_{\psi \in k^\vee} \bigoplus_{\chi \in \mu_{q+1}(k_2)^\vee} H_c^1(\mathbb{G}_m, \mathcal{L}_\psi \otimes \mathcal{K}_\chi) \simeq \bigoplus_{\psi \neq 1 \in k^\vee} \bigoplus_{\chi \neq 1 \in \mu_{q+1}(k_2)^\vee} H_c^1(\mathbb{G}_m, \mathcal{L}_\psi \otimes \mathcal{K}_\chi) \oplus \bigoplus_{\psi \in k^\vee} \psi \tag{7.3}$$

as $\overline{\mathbb{Q}}_\ell[k \times \mu_{q+1}(k_2)]$-modules. By (7.1), (7.2) and (7.3), the required assertion follows. ☐

For a character $\psi \in k^\vee$ and an element $\zeta \in k^\times$, we denote by ψ_ζ the character $x \mapsto \psi(\zeta x)$. We consider a character group $(k^\times)^\vee$ as a subgroup of $(k_2^\times)^\vee$ by $\mathrm{Nr}_{k_2/k}^\vee$.

Lemma 7.2. *We have an isomorphism* $H^1(X_{\mathrm{DL}}, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\tilde{\chi} \in (k_2^\times)^\vee \backslash (k^\times)^\vee} \tilde{\chi}$ *as* $\overline{\mathbb{Q}}_\ell[k_2^\times]$*-modules.*

Proof. By Lemma 7.1, we take a basis $\{e_{\psi,\chi}\}_{\psi \in k^\vee \backslash \{0\}, \chi \in \mu_{q+1}(k_2)^\vee \backslash \{1\}}$ of $H^1(X_{\mathrm{DL}}, \overline{\mathbb{Q}}_\ell)$ over $\overline{\mathbb{Q}}_\ell$ such that $k \times \mu_{q+1}(k_2)$ acts on $e_{\psi,\chi}$ by $\psi \times \chi$. For $\zeta \in k_2^\times$ and $a \in k$, we have $\beta_\zeta \circ \alpha_a \circ \beta_\zeta^{-1} = \alpha_{\zeta^{q+1} a}$ in $\mathrm{Aut}_{k_2}(X_{\mathrm{DL}})$. Hence, $\zeta \in k_2^\times$ acts on $H^1(X_{\mathrm{DL}}, \overline{\mathbb{Q}}_\ell)$ by $\zeta \colon e_{\psi,\chi} \mapsto c_{\psi,\chi,\zeta} e_{\psi_{\zeta^{-(q+1)}},\chi}$ with some constant $c_{\psi,\chi,\zeta} \in \overline{\mathbb{Q}}_\ell^\times$. Therefore, we acquire an isomorphism $H^1(X_{\mathrm{DL}}, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\chi \in \mu_{q+1}(k_2)^\vee \backslash \{1\}} \mathrm{Ind}_{\mu_{q+1}(k_2)}^{k_2^\times}(\chi)$ as $\overline{\mathbb{Q}}_\ell[k_2^\times]$-modules. Hence, th required assertion follows. ☐

Proposition 7.3. *We have isomorphisms*

$$H^1(\overline{\mathbf{Y}}_{1,2}^{\mathrm{c}}, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\tilde{\chi} \in (k_2^\times)^\vee \backslash (k^\times)^\vee} (\tilde{\chi} \circ \lambda) \otimes (\tilde{\chi}^q \circ \kappa_1),\ H^1(\overline{\mathbf{Y}}_{2,1}^{\mathrm{c}}, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\tilde{\chi} \in (k_2^\times)^\vee \backslash (k^\times)^\vee} (\tilde{\chi} \circ \lambda) \otimes (\tilde{\chi} \circ \kappa_1)$$

as $(I_K \times \mathcal{O}_D^\times)$*-representations over* $\overline{\mathbb{Q}}_\ell$*.*

Proof. This follows from Lemma 5.1, Lemma 6.2 and Lemma 7.2. ☐

Let X_{AS} be the smooth compactification of an affine curve X'_{AS} over k defined by $z^q - z = w^2$. Then $a \in k$ acts on X_{AS} by $(z, w) \mapsto (z + a, w)$. By this action, we consider $H^1(X_{\mathrm{AS}}, \overline{\mathbb{Q}}_\ell)$ as a $\overline{\mathbb{Q}}_\ell[k]$-module.

Lemma 7.4. *We assume that* q *is odd. Let* G *be a Galois group of a Galois extension* F *of* $k((s))$ *defined by* $z^q - z = 1/s^2$*. Let* G^r *be the upper numbering ramification filtration of* G*. Then* $G^r = G$ *if* $r \leq 2$*, and* $G^r = 1$ *if* $r > 2$*.*

Proof. We take $a \in F$ such that $a^q - a = 1/s^2$. Then $sa^{(q-1)/2}$ is a uniformizer of F. Let v_F be the normalized valuation of F. For $\sigma \in G$ and an integer i, the condition $v_F(\sigma(sa^{(q-1)/2}) - sa^{(q-1)/2}) \geq i$ is equivalent to the condition $v_F(\sigma(a) - a) \geq i - 3$. Hence, the claim follows. $\square$

Lemma 7.5. *We assume that q is odd. Then we have $H^1(X_{\mathrm{AS}}, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\psi \in k^\vee \setminus \{1\}} \psi$ as $\overline{\mathbb{Q}}_\ell[k]$-modules.*

Proof. We have $H^1(X_{\mathrm{AS}}, \overline{\mathbb{Q}}_\ell) \simeq H^1_c(X'_{\mathrm{AS}}, \overline{\mathbb{Q}}_\ell)$, because the complement $X_{\mathrm{AS}} \setminus X'_{\mathrm{AS}}$ consists of one point. The curve X'_{AS} is a finite etale Galois covering of $\mathbb{A}^1$ with a Galois group k by $(z, w) \mapsto w$. For $\psi \in k^\vee$, let $\mathcal{L}_{2,\psi}$ be the smooth $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{A}^1$ defined by the covering X'_{AS} and ψ. Then we have $H^1_c(X'_{\mathrm{AS}}, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\psi \in k^\vee \setminus \{1\}} H^1_c(\mathbb{A}^1, \mathcal{L}_{2,\psi})$ as $\overline{\mathbb{Q}}_\ell[k]$-modules. By Lemma 7.4 and the Grothendieck-Ogg-Shafarevich formula, we have $\dim H^1_c(\mathbb{A}^1, \mathcal{L}_{2,\psi}) = 1$ and $H^1_c(\mathbb{A}^1, \mathcal{L}_{2,\psi}) \simeq \psi$ as $\overline{\mathbb{Q}}_\ell[k]$-modules for $\psi \in k^\vee \setminus \{1\}$. Hence, the assertion follows. $\square$

We put $U_D = \{d \in \mathcal{O}_D^\times \mid \kappa_1(d) \in k^\times\}$. If q is odd, we put

$$\tau_{\chi,\psi} = \mathrm{Ind}_{I_L}^{I_K} \big((\chi \circ \lambda_1^{q+1}) \otimes (\psi^2 \circ \mathrm{Tr}_{k_2/k} \circ \lambda_2) \big), \ \theta_{\chi,\psi} = (\chi \circ \kappa_1) \otimes (\psi \circ \mathrm{Tr}_{k_2/k} \circ \kappa_2)$$

and $\rho_{\chi,\psi} = \mathrm{Ind}_{U_D}^{\mathcal{O}_D^\times} \theta_{\chi,\psi}$ for $\chi \in (k^\times)^\vee$ and $\psi \in k^\vee \setminus \{1\}$. We note that $\dim \rho_{\chi,\psi} = q + 1$.

Proposition 7.6. *We assume that q is odd. Then we have an isomorphism*

$$\bigoplus_{\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})} H^1(\overline{\mathbf{X}}^{\mathrm{c}}_\zeta, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\chi \in (k^\times)^\vee} \bigoplus_{\psi \in k^\vee \setminus \{1\}} \tau_{\chi,\psi} \otimes \rho_{\chi,\psi}$$

as representations of $I_K \times \mathcal{O}_D^\times$.

Proof. The actions of I_L and U_D on $\bigoplus_{\zeta \in k^\times} H^1(\overline{\mathbf{X}}^{\mathrm{c}}_\zeta, \overline{\mathbb{Q}}_\ell)$ factor through $k^\times \times k$ by Proposition 5.4 and Corollary 6.5. On the other hand, the action of $k^\times \times k$ on $\bigoplus_{\zeta \in k^\times} H^1(\overline{\mathbf{X}}^{\mathrm{c}}_\zeta, \overline{\mathbb{Q}}_\ell)$ is induced from an action of $\{1\} \times k$ on $H^1(\overline{\mathbf{X}}^{\mathrm{c}}_1, \overline{\mathbb{Q}}_\ell)$. Hence, we have $\bigoplus_{\zeta \in k^\times} H^1(\overline{\mathbf{X}}^{\mathrm{c}}_\zeta, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\chi \in (k^\times)^\vee} \bigoplus_{\psi \in k^\vee \setminus \{1\}} \chi \otimes \psi$ as representations of $k^\times \times k$ by Lemma 7.5. Therefore, we have an isomorphism

$$\bigoplus_{\zeta \in k^\times} H^1(\overline{\mathbf{X}}^{\mathrm{c}}_\zeta, \overline{\mathbb{Q}}_\ell) \simeq \bigoplus_{\chi \in (k^\times)^\vee} \bigoplus_{\psi \in k^\vee \setminus \{1\}} (\chi \circ \lambda_1^{q+1}) \otimes (\psi^2 \circ \mathrm{Tr}_{k_2/k} \circ \lambda_2) \otimes \theta_{\chi,\psi}$$

as representations of $I_L \times U_D$ by Proposition 5.4 and Corollary 6.5. Inducing this representation from I_L to I_K and U_D to $\mathcal{O}_D^\times$, we obtain the isomorphism in the assertion. $\square$

Let $Z \subset Q$ be a subgroup consisting of $g(1, 0, \gamma)$ with $\gamma^2 + \gamma = 0$, and ϕ be the unique non-trivial character of Z. By [BH, Lemma 22.2], there exists a unique irreducible two-dimensional representation τ of Q such that

$$\tau|_Z \simeq \phi^{\oplus 2}, \ \mathrm{Tr}\, \tau(g(\alpha, 0, 0)) = -1 \tag{7.4}$$

for $\alpha \in \mathbb{F}_4^\times \setminus \{1\}$. Then, it is easily checked that the determinant character of τ is trivial. Note that every two-dimensional irreducible representation of Q has a form $\tau \otimes \chi$ with $\chi \in (\mathbb{F}_4^\times)^\vee$, where we consider χ as a character of Q by $g(\alpha, \beta, \gamma) \mapsto \chi(\alpha)$.

Lemma 7.7. *The Q-representation $H^1(E, \overline{\mathbb{Q}}_\ell)$ is isomorphic to τ.*

Proof. The Q-representation $H^1(E, \overline{\mathbb{Q}}_\ell)$ satisfies (7.4) by Lemma 7.1. Hence, the assertion follows. $\square$

Let $\tau_{\zeta'}$ be the representation of W_K induced from the $(Q \rtimes \mathbb{Z})$-representation $H^1(E, \overline{\mathbb{Q}}_\ell)$ by $\Theta_{\zeta'}$. Then the restriction to I_K of $\tau_{\zeta'}$ is isomorphic to a representation induced from τ by Lemma 7.7.

We say that a continuous two-dimensional irreducible representation V of W_K over $\overline{\mathbb{Q}}_\ell$ is primitive, if there is no pair of a quadratic extension K' and a continuous character χ of $W_{K'}$ such that $V \simeq \mathrm{Ind}_{W_{K'}}^{W_K} \chi$.

Lemma 7.8. *The representation $\tau_{\zeta'}$ is primitive of Artin conductor 3.*

Proof. We use the notations in the proof of Lemma 6.11. The element $1/(\varpi^{1/3}\theta^3)$ is a uniformizer of $K^{\mathrm{ur}}(\varpi^{1/3},\theta)$. For $\sigma \in I_K$, we can show that

$$v\left(\sigma\left(\frac{1}{\varpi^{\frac{1}{3}}\theta^3}\right)-\frac{1}{\varpi^{\frac{1}{3}}\theta^3}\right)=\begin{cases}\frac{1}{24} & \text{if } \zeta_{3,\sigma}\neq 1,\\ \frac{1}{12} & \text{if } \zeta_{3,\sigma}=1,\ \nu_\sigma\neq 0,\\ \frac{1}{6} & \text{if } \zeta_{3,\sigma}=1,\ \nu_\sigma=0,\ \mu_\sigma\neq 0,\end{cases}$$

using $\sigma(\theta)\equiv\theta_\sigma \pmod{2/3}$. The claim on the Artin conductor follows from this.

The unique index 2 subgroup of $Q\rtimes\mathbb{Z}$ is $Q\rtimes 2\mathbb{Z}$, because Q has no index 2 subgroup. Hence, if $\tau_{\zeta'}$ is not primitive, it is induced from a character of W_{K_2}. However, this is impossible, because the restriction of $\tau_{\zeta'}$ to W_{K_2} is irreducible. □

We define a character $\lambda_\xi : W_K \to k^\times$ by $\lambda_\xi(\sigma)=\bar{\xi}_\sigma$. For $\zeta'\in k^\times$ and $\chi\in(k^\times)^\vee$, we put

$$\tau_{\zeta',\chi}=\tau_{\zeta'}\otimes(\chi\circ\lambda_\xi),\ \theta_{\zeta',\chi}=(\chi\circ\kappa_1)\otimes(\phi\circ\mathrm{Tr}_{k_2/\mathbb{F}_2}(\zeta'^{-2}\kappa_2)),\ \rho_{\zeta',\chi}=\mathrm{Ind}_{U_D}^{\mathcal{O}_D^\times}\theta_{\zeta',\chi}.$$

In the sequel, we consider $\tau_{\zeta',\chi}$ as a representation of I_K.

Proposition 7.9. *We assume that q is even. Let $\zeta'\in k^\times$. Then we have an isomorphism*

$$\bigoplus_{\zeta\in k_2^\times} H^1(\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta,\zeta'},\overline{\mathbb{Q}}_\ell)\simeq\bigoplus_{\chi\in(k^\times)^\vee}\tau_{\zeta',\chi}\otimes\rho_{\zeta',\chi}$$

as representations of $I_K\times\mathcal{O}_D^\times$.

Proof. The actions of I_K and U_D on $\bigoplus_{\zeta\in k^\times}H^1(\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta,\zeta'},\overline{\mathbb{Q}}_\ell)$ factor through $Q\times k^\times$ by Proposition 5.4 and Proposition 6.12. On the other hand, the action of $Q\times k^\times$ on $\bigoplus_{\zeta\in k^\times}H^1(\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta,\zeta'},\overline{\mathbb{Q}}_\ell)$ is induced from an action of Q on $H^1(\overline{\mathbf{X}}^{\mathrm{c}}_{1,\zeta'},\overline{\mathbb{Q}}_\ell)$. Hence, we have $\bigoplus_{\zeta\in k^\times}H^1(\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta,\zeta'},\overline{\mathbb{Q}}_\ell)\simeq\bigoplus_{\chi\in(k^\times)^\vee}\tau\otimes\chi$ as representations of $Q\times k^\times$. Therefore, we have an isomorphism

$$\bigoplus_{\zeta\in k^\times} H^1(\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta,\zeta'},\overline{\mathbb{Q}}_\ell)\simeq\bigoplus_{\chi\in(k^\times)^\vee}\tau_{\zeta',\chi}\otimes\theta_{\zeta',\chi}$$

as representations of $I_K\times U_D$ by Proposition 5.4 and Proposition 6.12. Inducing this representation from U_D to $\mathcal{O}_D^\times$, we obtain the isomorphism in the assertion. □

7.2 Genus calculation

Lemma 7.10. *We have* $\dim H^1_{\mathrm{c}}(\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}},\overline{\mathbb{Q}}_\ell)=2q^3-2q+1$.

Proof. It suffices to show $\dim H^1_{\mathrm{c}}\bigl((\mathrm{LT}_1(\mathfrak{p}^3)/\varpi^{\mathbb{Z}})_{\mathbf{C}},\overline{\mathbb{Q}}_\ell\bigr)=4q^3-4q+2$, because

$$\dim H^1_{\mathrm{c}}\bigl((\mathrm{LT}_1(\mathfrak{p}^3)/\varpi^{\mathbb{Z}})_{\mathbf{C}},\overline{\mathbb{Q}}_\ell\bigr)=2\dim H^1_{\mathrm{c}}(\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}},\overline{\mathbb{Q}}_\ell).$$

For an irreducible smooth representation π of $GL_2(K)$, we write $c(\pi)$ for the conductor of π. By Proposition 2.1, we have

$$H^1_{\mathrm{c}}\bigl((\mathrm{LT}_1(\mathfrak{p}^3)/\varpi^{\mathbb{Z}})_{\mathbf{C}},\overline{\mathbb{Q}}_\ell\bigr)\simeq\bigoplus_{\pi}\bigl(\pi^{K_1(\mathfrak{p}^3)}\bigr)^{\oplus 2\dim\mathrm{LJ}(\pi)}\oplus\bigoplus_{\chi}(\mathrm{St}\otimes\chi)^{K_1(\mathfrak{p}^3)},$$

where π runs through irreducible cuspidal representations of $GL_2(K)$ such that $c(\pi)\leq 3$ and the central character of π is trivial on ϖ, and χ runs through characters of $K^\times$ such that $c(\mathrm{St}\otimes\chi)\leq 3$ and $\chi(\varpi^2)=1$. We have the following list of discrete series representations π of $GL_2(K)$ such that $c(\pi)\leq 3$ and the central character of π is trivial on ϖ.

(1) $\pi\simeq\mathrm{St}\otimes\chi$ for an unramified character $\chi\colon K^\times\to\overline{\mathbb{Q}}_\ell^\times$ such that $\chi(\varpi^2)=1$. Then $c(\pi)=1$ and $\dim\mathrm{LJ}(\pi)=1$. There are two such representations.

(2) $\pi\simeq\mathrm{St}\otimes\chi$ for a tamely ramified character $\chi\colon K^\times\to\overline{\mathbb{Q}}_\ell^\times$ that is not unramified and satisfies $\chi(\varpi^2)=1$. Then $c(\pi)=2$ and $\dim\mathrm{LJ}(\pi)=1$. There are $2(q-2)$ such representations.

(3) $\pi \simeq \pi_\chi$, in the notation of [BH, 19.1], for a character $\chi\colon K_2^\times \to \overline{\mathbb{Q}}_\ell^\times$ of level zero such that χ does not factor through $\mathrm{Nr}_{K_2/K}$ and $\chi(\varpi) = 1$. Then $c(\pi) = 2$ and $\dim \mathrm{LJ}(\pi) = 2$. There are $q(q-1)/2$ such representations.

(4) $\pi \simeq \pi_\chi$, in the notation of [BH, 19.3], for a character $\chi\colon L^\times \to \overline{\mathbb{Q}}_\ell^\times$ of level one such that $\chi(\varpi) = 1$. Then $c(\pi) = 3$ and $\dim \mathrm{LJ}(\pi) = q+1$. There are $2(q-1)^2$ such representations.

We note that $\dim \pi^{K_1(\mathfrak{p}^3)} = 4 - c(\pi)$ if π is a discrete series representation of $GL_2(K)$ such that $c(\pi) \leq 3$. Then we obtain the claim by taking a summation according to the above list. $\square$

For an affinoid rigid space X, a Zariski subaffinoid of X is the inverse image of a nonempty open subscheme of $\overline{X}$ under the reduction map $X \to \overline{X}$.

Proposition 7.11. *Let W be a wide open rigid curve over a finite extension of $\widehat{K}^{\mathrm{ur}}$ with a stable covering $\{(U_i, U_i^{\mathrm{u}})\}_{i \in I}$. Let X be a subaffinoid space of W such that $\overline{X}$ is a connected smooth curve with a positive genus. Then there exists $i \in I$ such that X is a Zariski subaffinoid of U_i^{u}.*

Proof. Assume that $X \cap U_i^{\mathrm{u}}$ is contained in a finite union of residue classes of X for any $i \in I$. Then a Zariski subaffinoid of X appears in an open annulus. This is a contradiction, because $\overline{X}$ has a positive genus. Hence there exists $i' \in I$ such that $X \cap U_{i'}^{\mathrm{u}}$ is not contained in any finite union of residue class of X. We fix such i' in the sequel.

Then some open irreducible subscheme of the reduction of $X \cap U_{i'}^{\mathrm{u}}$ does not go to one point in $\overline{X}$ under the natural map $\overline{X \cap U_{i'}^{\mathrm{u}}} \to \overline{X}$. Let Y be the inverse image of such an open subscheme under the reduction map $X \cap U_{i'}^{\mathrm{u}} \to \overline{X \cap U_{i'}^{\mathrm{u}}}$. Then we see that Y is a Zariski subaffinoid of X by [CM, Lemma 2.24 (i)]. Each connected component of $X \setminus Y$ is an open disk, and included in $U_{i'}^{\mathrm{u}}$ or U_i^{u} for $i \neq i'$ or an open annulus outside the underlying affinoids. This can be checked by applying [CM, Corollary 2.39] to every closed disk in a connected component of $X \setminus Y$. Hence, $X \cap U_{i'}^{\mathrm{u}}$ is a Zariski subaffinoid of X. If $X \cap U_{i'}^{\mathrm{u}} \neq X$, then $U_{i'}^{\mathrm{u}}$ is connected to an open disk in U_i^{u} for $i \neq i'$ or in an open annulus outside the underlying affinoids. This is a contradiction. Therefore, we have $X \subset U_i^{\mathrm{u}}$. Then we obtain the claim by [CM, Lemma 2.24 (i)]. $\square$

Lemma 7.12. *Let W be a wide open rigid curve over a finite extension of $\widehat{K}^{\mathrm{ur}}$ with a stable covering. Let X be a subaffinoid space of W such that $\overline{X}$ is a connected smooth curve with genus zero. Then there is a basic wide open subspace of W such that its underlying affinoid is X.*

Proof. We note that we have the claim if X appears in an open subannulus of W. Let $\{(U_i, U_i^{\mathrm{u}})\}_{i \in I}$ be the stable covering of W.

First, we consider the case where $X \cap U_i^{\mathrm{u}}$ is contained in a finite union of residue classes of X for any $i \in I$. Then a Zariski subaffinoid of X appears in an open annulus. Further, X itself appears in the open annulus, because X is connected. Hence, we have the claim in this case.

Therefore, we may assume that there exists $i' \in I$ such that $X \cap U_{i'}^{\mathrm{u}}$ is not contained in any finite union of residue class of X. We fix such i'. By the same argument as in the proof of Proposition 7.11, we have $X \subset U_i^{\mathrm{u}}$. If the image of the induced map $\overline{X} \to \overline{U}_i^{\mathrm{u}}$ is one point, we have the claim because X appears in an open disk. Otherwise, X is a Zariski subaffinoid of U_i^{u}, and we have the claim. $\square$

We consider the natural level-lowering map

$$\pi_f\colon \mathbf{X}_1(\mathfrak{p}^3) \to \mathbf{X}_1(\mathfrak{p}^2);\ (u, X_3) \mapsto (u, X_2).$$

Lemma 7.13. *The connected components of $\mathbf{W}_{1,2'}$, $\mathbf{W}_{1,3'}$, $\mathbf{W}_{2,1'}$ and $\mathbf{W}_{4,1'} \cup \mathbf{W}_{5,1'} \cup \mathbf{W}_{6,1'}$ are not open balls.*

Proof. Let $\mathbf{W}_0'$ be a subannulus of $\mathbf{W}_0$ defined by $v(u) < 1/(q(q+1))$. Then we have $\pi_f^{-1}(\mathbf{W}_{k^\times}) = \mathbf{W}_{2,1'}$, $\pi_f^{-1}(\mathbf{W}_\infty) = \mathbf{W}_{4,1'} \cup \mathbf{W}_{5,1'} \cup \mathbf{W}_{6,1'}$ and $\pi_f^{-1}(\mathbf{W}_0') = \mathbf{W}_{1,2'} \cup \mathbf{W}_{1,3'}$. Hence we have the claim by Proposition 3.1 and [Co, Lemma 1.4]. $\square$

The smooth projective curves $\overline{\mathbf{Y}}_{1,2}^{\mathrm{c}}$ and $\overline{\mathbf{Y}}_{2,1}^{\mathrm{c}}$ have defining equations $X^qY - XY^q = Z^{q+1}$ determined by the equation in Proposition 4.2 and Proposition 4.3. The infinity points of $\overline{\mathbf{Y}}_{1,2}$ consist of $P_a^+ = (a, 1, 0)$ for $a \in k$ and $P_\infty^+ = (1, 0, 0)$. The infinity points of $\overline{\mathbf{Y}}_{2,1}$ consist of $P_a^- = (a, 1, 0)$ for $a \in k$ and $P_\infty^- = (1, 0, 0)$.

For a wide open space W, let $e(W)$ be the number of the ends of W, and $g(W)$ be the genus of W (cf. [CM, p. 369 and p. 380]). For a proper smooth curve C, we write $g(C)$ for the genus of C.

Theorem 7.14. *The covering $\mathcal{C}_1(\mathfrak{p}^3)$ is a semi-stable covering of $\mathbf{X}_1(\mathfrak{p}^3)$ over some finite extension.*

Proof. We consider the stable covering of $\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}}$ by Proposition 4.11. Then $\overline{\mathbf{Y}}^{\mathrm{c}}_{1,2}$ and $\overline{\mathbf{Y}}^{\mathrm{c}}_{2,1}$ appear in the stable reduction of $\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}}$ by Proposition 7.11. The point P_0^+ is the unique infinity point of $\overline{\mathbf{Y}}_{1,2}$ whose tube is contained in $\mathbf{W}^+_{1,1'}$, because $v(X_3) > 1/(q^3(q^2-1))$ in $\mathbf{W}^+_{1,1'}$. Similarly, P_0^- is the unique infinity point of $\overline{\mathbf{Y}}_{2,1}$ whose tube is contained in $\mathbf{W}^-_{1,1'}$. Hence, we have $e(\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}}) \geq 2q$ by Lemma 7.13. Therefore, we have $g(\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}}) \leq q^3 - 2q + 1$ by Lemma 7.10. On the other hand, we have

$$g(\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}}) \geq g(\overline{\mathbf{Y}}^{\mathrm{c}}_{1,2}) + g(\overline{\mathbf{Y}}^{\mathrm{c}}_{2,1}) + \begin{cases} \sum_{\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})} g(\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta}) & \text{if } q \text{ is odd,} \\ \sum_{\zeta \in k_2^{\times},\ \zeta' \in k^{\times}} g(\overline{\mathbf{X}}^{\mathrm{c}}_{\zeta,\zeta'}) & \text{if } q \text{ is even,} \end{cases}$$

where the summation on the right hand side is $q^3 - 2q + 1$ by Proposition 7.3, Proposition 7.6 and Proposition 7.9. Then the affinoids $\mathbf{Y}_{1,2}$, $\mathbf{Y}_{2,1}$, $\mathbf{X}_{\zeta}$ for $\zeta \in \mu_{2(q^2-1)}(k^{\mathrm{ac}})$ and $\mathbf{X}_{\zeta,\zeta'}$ for $\zeta \in k_2^{\times}$ and $\zeta' \in k^{\times}$ are underlying affinoids of basic wide open spaces in the stable covering by Proposition 7.11 and Lemma 7.13. Therefore, by the above genus inequalities, we see that $e(\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}}) = 2q$ and the connected components of $\mathbf{W}_{1,2'}$, $\mathbf{W}_{1,3'}$, $\mathbf{W}_{2,1'}$ and $\mathbf{W}_{4,1'} \cup \mathbf{W}_{5,1'} \cup \mathbf{W}_{6,1'}$ are open annuli.

The connected components of $\mathbf{X}_1(\mathfrak{p}^3) \setminus \mathbf{Z}^0_{1,1}$ are two wide open spaces, because each connected component is connected to $\mathbf{Z}^0_{1,1}$ at an open subannulus by Lemma 7.12. Then we see that these two wide open spaces are basic wide open spaces with underlying affinoids $\mathbf{Y}_{1,2}$ and $\mathbf{Y}_{2,1}$ by the above genus inequalities. Therefore we have the claim by Proposition 4.7, Proposition 4.9 and Proposition 4.10. ☐

7.3 Structure of cohomology

In this subsection, we study the action of $I_K \times \mathcal{O}_D^{\times}$ on ℓ-adic cohomology of $\mathbf{X}_1(\mathfrak{p}^3)$. We put $(W_K \times D^{\times})^0 = \{(\sigma, \varphi^{-r_\sigma}) \in W_K \times D^{\times}\}$. Although it is possible to study the action of $(W_K \times D^{\times})^0$ using the result of Section 6, here we keep us to study the inertia action for simplicity. The result in this subsection is essentially used in [IT2].

Let $\mathcal{X}_1(\mathfrak{p}^3)$ be a semi-stable formal scheme constructed from $\mathcal{C}_1(\mathfrak{p}^3)$ by [IT1, Theorem 3.5]. The semi-stable reduction of $\mathcal{X}_1(\mathfrak{p}^3)$ means the underlying reduced scheme of $\mathcal{X}_1(\mathfrak{p}^3)$, which is denoted by $\mathcal{X}_1(\mathfrak{p}^3)_{k^{\mathrm{ac}}}$.

Lemma 7.15. *The smooth projective curves $\overline{\mathbf{Y}}^{\mathrm{c}}_{1,2}$ and $\overline{\mathbf{Y}}^{\mathrm{c}}_{2,1}$ intersect with $\overline{\mathbf{Z}}^{\mathrm{c}}_{1,1}$ at P_0^+ and P_0^- respectively in the stable reduction $\mathcal{X}_1(\mathfrak{p}^3)_{k^{\mathrm{ac}}}$.*

Proof. We see this from the proof of Theorem 7.14. ☐

Let Γ be the graph defined by the following:

- The set of the vertices of Γ consists of P_0, P_∞, P_a^+ and P_a^- for $a \in \mathbb{P}^1(k) \setminus \{0\}$.
- The set of the edges of Γ consists of $P_0P_a^+$, $P_0P_a^-$, $P_\infty P_a^+$ and $P_\infty P_a^-$ for $a \in \mathbb{P}^1(k) \setminus \{0\}$.

We note that P_a^+ and P_a^- for $a \in \mathbb{P}^1(k) \setminus \{0\}$ are points of $\overline{\mathbf{Y}}^{\mathrm{c}}_{1,2}$ and $\overline{\mathbf{Y}}^{\mathrm{c}}_{2,1}$ that are not on $\overline{\mathbf{Z}}^{\mathrm{c}}_{1,1}$ by Lemma 7.15. Let $H^1(\Gamma, \overline{\mathbb{Q}}_\ell)$ be the cohomology group of Γ with coefficients in $\overline{\mathbb{Q}}_\ell$ (cf. [IT1, Section 2]). The group $I_K \times \mathcal{O}_D^{\times}$ acts on P_a^+ and P_a^- for $a \in \mathbb{P}^1(k) \setminus \{0\}$ via the action on $\overline{\mathbf{Y}}^{\mathrm{c}}_{1,2}$ and $\overline{\mathbf{Y}}^{\mathrm{c}}_{2,1}$. Let $I_K \times \mathcal{O}_D^{\times}$ act on P_0 and P_∞ trivially. By this action, we consider $H^1(\Gamma, \overline{\mathbb{Q}}_\ell)$ as a $\overline{\mathbb{Q}}_\ell[I_K \times \mathcal{O}_D^{\times}]$-module.

Theorem 7.16. *We have an exact sequence*

$$0 \longrightarrow H^1(\Gamma, \overline{\mathbb{Q}}_\ell) \longrightarrow H^1_{\mathrm{c}}(\mathbf{X}_1(\mathfrak{p}^3)_{\mathbf{C}}, \overline{\mathbb{Q}}_\ell) \longrightarrow H^1(\mathcal{X}_1(\mathfrak{p}^3)_{k^{\mathrm{ac}}}, \overline{\mathbb{Q}}_\ell)^*(-1) \longrightarrow 0$$

as representations of $(W_K \times D^{\times})^0$. Further, as $(I_K \times \mathcal{O}_D^{\times})$-representations, $H^1(\mathcal{X}_1(\mathfrak{p}^3)_{k^{\mathrm{ac}}}, \overline{\mathbb{Q}}_\ell)$ is isomorphic to

$$\bigoplus_{\tilde{\chi} \in (k_2^{\times})^{\vee} \setminus (k^{\times})^{\vee}} \big((\tilde{\chi} \circ \lambda) \otimes (\tilde{\chi} \circ \kappa_1 \oplus \tilde{\chi} \circ \kappa_1)\big) \oplus \begin{cases} \bigoplus_{\chi \in (k^{\times})^{\vee}} \bigoplus_{\psi \in k^{\vee} \setminus \{1\}} \tau_{\chi,\psi} \otimes \rho_{\chi,\psi} & \textit{if } q \textit{ is odd,} \\ \bigoplus_{\zeta' \in k^{\times}} \bigoplus_{\chi \in (k^{\times})^{\vee}} \tau_{\zeta',\chi} \otimes \rho_{\zeta',\chi} & \textit{if } q \textit{ is even,} \end{cases}$$

and $H^1(\Gamma, \overline{\mathbb{Q}}_\ell)$ is isomorphic to

$$1 \oplus \bigoplus_{\chi \in (k^{\times})^{\vee}} \big((\chi \circ \lambda^{q+1}) \otimes (\chi \circ \kappa_1^{q+1})\big)^{\oplus 2}.$$

Proof. The existence of the exact sequence follows from [IT1, Theorem 5.3] and Lemma 7.15 using the Poincaré duality (cf. [Far1, Proposition 5.9.2]). We know the structure of $H^1(\mathcal{X}_1(\mathfrak{p}^3)_{k^{\mathrm{ac}}}, \overline{\mathbb{Q}}_\ell)$ by Proposition 7.3, Proposition 7.6 and Proposition 7.9.

We study the structure of $H^1(\Gamma, \overline{\mathbb{Q}}_\ell)$. By Lemma 5.1 and Lemma 6.2, the action of $I_K \times \mathcal{O}_D^\times$ on $H^1(\Gamma, \overline{\mathbb{Q}}_\ell)$ factor through $k^\times$. We can check that $H^1(\Gamma, \overline{\mathbb{Q}}_\ell) \simeq 1 \oplus \bigoplus_{\chi \in (k^\times)^\vee} \chi^{\oplus 2}$ as representations of $k^\times$. Hence, the claim follows from Lemma 5.1 and Lemma 6.2. □

References

[BH] C. J. Bushnell and G. Henniart, *The local Langlands conjecture for GL(2)*, Grundlehren der Mathematischen Wissenschaften, vol 335, Springer-Verlag, Berlin, 2006.

[Co] R. Coleman, *Stable maps of curves*, Kazuya Kato's fiftieth birthday. Doc. Math. 2003, Extra Vol., 217–225.

[CM] R. Coleman and K. McMurdy, *Stable reduction of* $X_0(p^3)$, Algebra and Number Theory **4** (2010), no. 4 357–431, with an appendix by Everette W. Howe.

[Da] J.-F. Dat, *Théorie de Lubin-Tate non-abélienne et représentations elliptiques*, Invent. Math. **169** (2007), no. 1, 75–152.

[De] P. Deligne, *Formes modulaires et représentations de* GL(2), Modular functions of one variable II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), pp. 55–105. Lecture Notes in Math., Vol. 349, Springer, Berlin, 1973.

[DR] P. Deligne and M. Rapoport, *Les Schémas de modules de courbes elliptiques*, Lecture Notes in Math. **349** (1973), 143–316.

[Fal] G. Faltings, *A relation between two moduli spaces studied by V. G. Drinfeld*, Algebraic number theory and algebraic geometry, 115–129, Contemp. Math., 300, Amer. Math. Soc., Providence, RI, 2002.

[Far1] L. Fargues, *Cohomologie des espaces de modules de groupes p-divisibles et correspondances de Langlands locales*, Astérisque No. 291 (2004), 1–199.

[Far2] L. Fargues, *Dualité de Poincaré et involution de Zelevinsky dans la cohomologie équivariante des espaces rigides*, preprint. Available at http://www-irma.u-strasbg.fr/~fargues/Prepublications.html

[FGL] L. Fargues, A. Genestier and V. Lafforgue, *L'isomorphisme entre les tours de Lubin-Tate et de Drinfeld*, Progress in Mathematics, 262. Birkhäuser Verlag, Basel, 2008.

[GH] B. Gross and M. Hopkins, *Equivariant vector bundles on the Lubin-Tate moduli space*, Topology and Representation Theory (Evanston, IL, 1992), Contemp. Math. **158** (1994), 23–88.

[Ha] M. Hazewinkel, *Formal groups and applications*, Academic Press, 1978.

[IT1] N. Imai and T. Tsushima, *Cohomology of rigid curves with semi-stable coverings*, preprint, arXiv:1109.2824.

[IT2] N. Imai and T. Tsushima, *Geometric realization of the local Langlands correspondence for representations of conductor three*, preprint, arXiv:1205.0734.

[IT3] N. Imai and T. Tsushima, *Explicit construction of semi-stable models of Lubin-Tate curves with low level.* Available at http://www.kurims.kyoto-u.ac.jp/~naoki/proceedings.html

[M1] Y. Mieda, *Non-cuspidality outside the middle degree of ℓ-adic cohomology of the Lubin-Tate tower*, Adv. Math. **225** (2010), no. 4, 2287–2297.

[M2] Y. Mieda, *Geometric approach to the local Jacquet-Langlands correspondence*, preprint, arXiv:1112.6116.

[Se] J.-P. Serre, *Représentations linéaires des groupes finis*, Hermann, Paris, 1967.

[SGA5] *Cohomologie l-adique et Fonctions L*, Séminaire de Géométrie Algébrique du Bois-Marie 1965–1966, dirigé par A. Grothendieck. Lecture Notes in Mathematics, Vol. 589, Springer-Verlag, Berlin-New York, 1977.

[St1] M. Strauch, *On the Jacquet-Langlands correspondence in the cohomology of the Lubin-Tate deformation tower*, Astérisque (2005) no. 298, 391–410, Formes Automorphics (I).

[St2] M. Strauch, *Deformation spaces of one-dimensional formal modules and their cohomology*, Adv. Math. **217** (2008), no. 3, 889–951.

[St3] M. Strauch, *Geometrically connected components of Lubin-Tate deformation spaces with level structures*, Pure Appl. Math. Q. 4 (2008), no. 4, Special Issue: In honor of Jean-Pierre Serre. Part 1, 1215–1232.

[We] A. Weil, *Exercices dyadiques*, Invent. Math. **27** (1974), 1–22.

Naoki Imai
Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502 Japan
naoki@kurims.kyoto-u.ac.jp

Takahiro Tsushima
Faculty of Mathematics, Kyushu University, 744 Motooka, Nishi-ku, Fukuoka city, Fukuoka 819-0395, Japan
tsushima@math.kyushu-u.ac.jp